arXiv:math/0103235v1 [math.CV] 30 Mar 2001

BIHOLOMORPHIC MAPPING ON THE BOUNDARY I

Won K. Park Department of Mathematics, University of Seoul wonkpark@hotmail.com

Abstract. We present a new proof of Chern-Ji's mapping theorem on a strongly pseudoconvex domain with differentiable spherical boundary. We show that a proper holomorphic self mapping of a strongly pseudoconvex domain with the real analytic boundary is biholomorphic.

0. Introduction and Preliminaries

We shall show that a bounded domain D is biholomorphic to an open ball B^{n+1} whenever the boundary bD is locally biholomorphic to the boundary of an open ball B^{n+1}.

Theorem 1. *Let D be a simply connected bounded domain in $\mathbb{C}^{n+1}$ with differentiable spherical boundary bD. Suppose that there is a biholomorphic mapping*

$$\phi \in H\left(U \cap D\right) \cap C^1\left(U \cap \overline{D}\right)$$

for a connected open neighborhood U of a point $p \in bD$ satisfying

$$\phi\left(U \cap bD\right) \subset bB^{n+1}.$$

Then the mapping ϕ is analytically continued to a biholomorphic mapping from D onto B^{n+1}.

Our result is a new proof of a weaker version of Chern-Ji's mapping theorem [CJ]. The main steps of our proof come as follows: We show that the inverse mapping ϕ^{-1} is analytically continued on the unit ball B^{n+1} to be a locally biholomorphic mapping

$$\varphi : B^{n+1} \to D.$$

We show that the mapping φ is a proper holomorphic mapping onto a universal covering Riemann domain over D. Thus the mapping φ is a biholomorphic mapping whenever D is simply connected.

We shall study on a proper holomorphic mapping ϕ between strongly pseudoconvex bounded domains D, D' with real analytic boundaries bD, bD'.

Theorem 2. *Let D, D' be strongly pseudoconvex bounded domains in $\mathbb{C}^{n+1}$ with real analytic boundaries bD, bD' and $\phi : D \to D'$ be a proper holomorphic mapping. Then the mapping ϕ is locally biholomorphic. If $D = D'$, then the mapping ϕ is a biholomorphic self mapping.*

Our result is a new proof of a weaker version of Pinchuk's mapping theorem [Pi]. The main steps of our proof come as follows: We show that the mapping ϕ is analytically continued along any path on bD as a locally biholomorphic mapping when bD is nonspherical so that the mapping $\phi : D \to D'$ are locally biholomorphic. From the study of Theorem 1, we show that the same is true when bD is spherical so that the mapping $\phi : D \to D'$ are locally biholomorphic. For the case of $D = D'$, we show that the boundary bD is necessarily spherical whenever the claim is not true. Then we show that there is a sequence of automorphisms $\phi_j \in Aut\left(D\right)$ and a sequence of points p_j on a compact subset $K \subset\subset D$ such that

$$\phi_j\left(p_j\right) \rightarrow bD$$

whenever the boundary bD is spherical and the claim is not true. We apply Wong-Rosay Theorem so that the domain D is biholomorphic to an open ball B^{n+1}. Then we obtain a contradiction that the mapping ϕ induces a nonautomorphic proper self mapping of an open ball B^{n+1} whenever the boundary bD is spherical and the claim is not true.

We remark that Theorem 1 is a weaker version of the following theorem:

Theorem 3 (cf. Chern-Ji [CJ). *] Let D be a simply connected bounded domain in $\mathbb{C}^{n+1}$ with continuous spherical boundary bD. Suppose that there is a biholomorphic mapping*

$$\phi \in H\left(U \cap D\right) \cap C\left(U \cap \overline{D}\right)$$

for a connected open neighborhood U of a point $p \in bD$ satisfying

$$\phi\left(U \cap bD\right) \subset bB^{n+1}.$$

Then the mapping ϕ is analytically continued to a biholomorphic mapping from D onto B^{n+1}.

We remark that Theorem 2 is a weaker version of the following theorem:

Theorem 4 (cf. Pinchuk [Pi). *] Let D, D' be strongly pseudoconvex bounded domains in $\mathbb{C}^{n+1}$ with the boundaries bD, bD' of class C^2 and $\phi : D \rightarrow D'$ be a proper holomorphic mapping. Then the mapping ϕ is locally biholomorphic. If $D = D'$, then the mapping ϕ is a biholomorphic self mapping.*

We presented parts of this article in the spring meeting of Korean Mathematical Society in 2000. This article is a preliminary version. Any comment shall be greatly appreciated. Send your comments to wonkpark@hotmail.com.

0.1. **Canonical normalizing mapping.**

Let M be a nondegenerate analytic real hypersurface in $\mathbb{C}^{n+1}$. For each point $p \in M$, there is a complex tangent hyperplane $H_p \subset T_pM$ so that there is a unit tangent vector $v_p \in T_pM$ perpendicular to the complex tangent hyperplane H_p with respect to the usual riemannian metric in $\mathbb{C}^{n+1} = \mathbb{R}^{2n+2}$. Then we can take a unique distinguished chain γ_p tangential to the direction v_p and passing through the complex tangent hyperplane H_p at the point p on M(cf. [Pa3]). Further, there is a distinguished normal parametrization on the chain γ_p having the same values up to order 2 of the straight real line to the direction v_p with the usual euclidean parametrization.. Therefore, we can take a distinguished normalizing mapping μ_p to Moser normal form

$$\mu_p : M \rightarrow \mu_p\left(M\right)$$

sending the germ M at the point p to a normal form such that $\mu_p(p)$ is the origin and $\mu_p\left(\gamma_p\right)$ is on the straightened chain of the normal form.

We take a local orientation near the point $p \in M$ so that the tangent vector v_p extends to a smooth unit vector field v on $M \cap U$ for an open neighborhood U of the point p such that $v_q \in T_qM$ is a unit tangent vector perpendicular to the complex tangent hyperplane H_q for each $q \in M \cap U$. Then we obtain a family of the distinguished normalizing mapping μ_q for $q \in M \cap U$:

$$\mu_q : M \to \mu_q (M)$$

associated the unit tangent vector $v_q \in T_q M \backslash H_q$. The distinguished normalizing mapping μ_q shall be called the canonical normalizing mapping associated Moser normal form.

Lemma 5. *Let $p_j \in M$ be a sequence of points converging to a point $p \in M$. Then there is a positive real number $\delta > 0$ such that*

1. *the mapping μ_{p_j} and its inverse $\mu_{p_j}^{-1}$ are analytically continued respectively on*

$$B(p_j; \delta) \quad \textit{and} \quad B(0; \delta)$$

as a biholomorphic mapping,

2. *the real hypersurface $\mu_{p_j}(M)$ is analytically continued on $B(0; \delta)$ by its defining equation,*
3. *the sequence $\mu_{p_j}^{-1}$ uniformly converges to μ_p^{-1} on $B(0; \delta)$ as a biholomorphic mapping,*
4. *the sequence $\mu_{p_j}(M)$ uniformly converges on $B(0; \delta)$ to the real hypersurface $\mu_p(M)$.*

Let H be the local automorphism group at the origin of the real hyperquadric

$$v = \langle z, z \rangle.$$

The isotropy subgroup $Aut_p(M)$ is naturally identified to the isotropy subgroup $Aut_0(\mu_p(M))$ by the following relation:

$$\mu_p \circ \phi \circ \mu_p^{-1} \in Aut_0(\mu_p(M)) \quad \text{for} \quad \phi \in Aut_p(M).$$

Note that every biholomorphic mapping φ between real hypersurfaces in normal form is faithfully represented by a natural group action of the isotropy subgroup H(cf. [Pa3]) such that

$$\varphi = N_e \quad \text{for} \quad e \in H.$$

Because $\mu_p(M)$ is in normal form, there is a natural identification of a local automorphism $\phi \in Aut_p(M)$ to an element

$$(U_\phi, a_\phi, \rho_\phi, r_\phi) \in H,$$

where $U_\phi, a_\phi, \rho_\phi, r_\phi$ are the normalizing parameters(cf. [Pa3]) of the mapping

$$\mu_p \circ \phi \circ \mu_p^{-1} \in Aut_0(\mu_p(M)).$$

Lemma 6. *If the isotropy subgroup $Aut_p(M)$ is compact, then $Aut_p(M)$ is isomorphic to the subgroup*

$$\{(U_\phi, a_\phi, \rho_\phi, r_\phi) \in H : \phi \in Aut_p(M)\}$$

as a Lie group.

0.2. **Preliminary Lemmas.** We have lemmas on the automorphism of B^{n+1}.

Lemma 7. *Let p, q be two distinct points on bB^{n+1} and $\phi \in Aut\left(bB^{n+1}\right)$ be a local automorphism of bB^{n+1} such that*

$$\phi(p) \neq q.$$

Then there is a unique decomposition

$$\phi = \psi \circ \varphi$$

where

$$\varphi \in Aut_p\left(bB^{n+1}\right), \quad \psi \in Aut_q\left(bB^{n+1}\right)$$

and the local automorphism ψ acts trivially on the complex tangent hyperplane of bB^{n+1} at the fixed point q.

Proof. Note that the isotropy subgroup $Aut_q\left(bB^{n+1}\right)$ acts on $bB^{n+1}\backslash q$ transitively. Further, there is a unique element $\psi \in Aut_q\left(bB^{n+1}\right)$ for each point displacement on $bB^{n+1}\backslash q$ by requiring the element ψ acts trivially on the complex tangent hyperplane of bB^{n+1} at the point q(cf. [Pa1]).

Let's put $p' = \phi(p)$. Since $p' \neq q$, we take a unique automorphism $\psi \in Aut_q\left(bB^{n+1}\right)$ such that

$$\psi(p') = p.$$

Then $\varphi \equiv \psi \circ \phi \in Aut_p\left(bB^{n+1}\right)$ so that

$$\phi = \psi^{-1} \circ \varphi.$$

This completes the proof. □

Lemma 8. *Let p, q be two distinct points on bB^{n+1} and $\phi \in Aut\left(bB^{n+1}\right)$ be a local automorphism of bB^{n+1} such that*

$$p' \equiv \phi(p) \neq q.$$

Then there is a unique decomposition

$$\phi = \varphi \circ \psi$$

where

$$\varphi \in Aut_{p'}\left(bB^{n+1}\right), \quad \psi \in Aut_q\left(bB^{n+1}\right)$$

and the local automorphism ψ acts trivially on the complex tangent hyperplane of bB^{n+1} at the fixed point q.

Proof. By Lemma 8, there is a decomposition

$$\phi^{-1} = \psi \circ \varphi$$

where

$$\varphi \in Aut_{p'}\left(bB^{n+1}\right), \quad \psi \in Aut_q\left(bB^{n+1}\right)$$

and the local automorphism ψ acts trivially on the complex tangent hyperplane of bB^{n+1} at the fixed point q. Hence we obtain

$$\phi = \varphi^{-1} \circ \psi^{-1}$$

where

$$\varphi^{-1} \in Aut_{p'}\left(bB^{n+1}\right), \quad \psi^{-1} \in Aut_{q}\left(bB^{n+1}\right).$$

This completes the proof. □

Lemma 9. *Let ϕ_j be a sequence of automorphisms of B^{n+1}. Suppose that the sequence ϕ_j converges to a holomorphic mapping λ uniformly on every compact subset of B^{n+1}. Then the mapping λ is either a constant mapping or an automorphism of B^{n+1}.*

Proof. Note that the mapping λ satisfies

$$\lambda\left(B^{n+1}\right) \subset \overline{B^{n+1}} \tag{1}$$

and a complex line is mapped to a complex line under the biholomorphic automorphism of the unit ball B^{n+1}.

Suppose that there is a complex line π such that

$$\pi \cap B^{n+1} \neq \emptyset \quad \text{and} \quad \lambda\left(\pi \cap B^{n+1}\right) = \emptyset.$$

Note that bB^{n+1} is strongly pseudoconvex so that, by the condition 1, there is a point $q \in bB^{n+1}$ satisfying

$$q = \lambda\left(\pi\right) \cap \overline{B^{n+1}}.$$

Then we obtain

$$\lambda|_{\pi \cap B^{n+1}} = q.$$

Let x be an interior point of $\pi \cap B^{n+1}$ and p' be an arbitrary point of B^{n+1} so that we take a complex line π' passes through x and p'. Since B^{n+1} is strongly pseudoconvex, the maximum modulus theorem of one complex variable yields

$$\lambda|_{\pi' \cap B^{n+1}} = q$$

so that the mapping λ is a constant mapping.

Suppose that the mapping λ is not a constant mapping. Then, for a complex line π satisfying

$$\pi \cap B^{n+1} \neq \emptyset,$$

there is a real number $\varepsilon > 0$ such that

$$\left|\phi_j\left(\pi \cap B^{n+1}\right)\right| \geq \varepsilon \tag{2}$$

where $\left|\phi_j\left(\pi \cap B^{n+1}\right)\right|$ is the area of the analytic disk

$$\phi_j\left(\pi \cap B^{n+1}\right).$$

We take a point $p \in \pi \cap bB^{n+1}$ so that

$$\phi_j\left(p\right) \to p' \in bB^{n+1}. \tag{3}$$

Then we take a point $p'' \in bB^{n+1}$ such that

$$p'' \notin \overline{\{\phi_j\left(p\right) : j \in \mathbb{N}^{+}\}},$$

if necessary, passing to a subsequence. We have the following decomposition

$$\phi_j = \varphi_j \circ \psi_j$$

where

$$\varphi_j \in Aut_{p'}\left(bB^{n+1}\right) \quad \text{and} \quad \psi_j \in Aut_{p''}\left(bB^{n+1}\right)$$

where the automorphisms ψ_j act trivially on the complex tangent hyperplane at the fixed point p''. Then we obtain

$$U_{\psi_j} = id_{n\times n}, \quad \rho_{\psi_j} = 1.$$

By the condition 3, there is a real number $e > 0$ such that

$$\left|a_{\psi_j}\right| \le e, \quad \left|r_{\psi_j}\right| \le e.$$

By the condition 2, there is a real number $e > 0$, if necessary, increasing e, such that

$$\left|a_{\varphi_j}\right| \le e.$$

Since the mapping λ is not a constant mapping, there is a real number $e > 0$, if necessary, increasing e, such that

$$e^{-1} \le \left|\rho_{\varphi_j}\right| \le e, \quad \left|r_{\varphi_j}\right| \le e.$$

Since bB^{n+1} is strongly pseudoconvex, we have

$$\left|U_{\varphi_j}\right| = 1.$$

Then the Jacobian determinant $\det \varphi_j'$ is uniformly bounded from the zero on an open neighborhood of the point $p \in bB^{n+1}$. By Hurwitz theorem, the mapping λ is locally biholomorphic and, further, the mapping λ is one-to-one. Hence the mapping λ is an automorphism of the unit ball B^{n+1}. This completes the proof. □

We have lemmas on the chain of bB^{n+1}.

Lemma 10. *Let $\gamma : [0,1] \to bB^{n+1}$ be a chain-segment on bB^{n+1}. Then there is a complex line π such that*

$$\gamma[0,1] \subset \pi \cap bB^{n+1}.$$

Proof. We take a point $p \in \gamma[0,1]$. Then the chain-segment $\mu_p\circ\gamma[0,1]$ on $\mu_p\left(bB^{n+1}\right)$ is on a complex line π'(cf. [Pa3]). Since the canonical normalizing mapping μ_p of the sphere bB^{n+1} is a fractional linear mapping(cf. [Pa3]), we take

$$\pi = \mu_p^{-1}\left(\pi'\right)$$

so that

$$\gamma[0,1] \subset \pi \cap bB^{n+1}.$$

This completes the proof. □

Lemma 11. *Let π be a complex line such that*

$$\pi \cap B^{n+1} \ne \emptyset.$$

Then the circle $\pi \cap bB^{n+1}$ is a chain on bB^{n+1}.

Proof. We take a point $p \in \pi \cap bB^{n+1}$. Then we obtain

$$\mu_p(\pi) \cap \mu_p\left(bB^{n+1}\right) = \mu_p\left(\pi \cap bB^{n+1}\right) \neq \emptyset.$$

Since the canonical normalizing mapping μ_p of the sphere bB^{n+1} is a fractional linear mapping(cf. [Pa3]), $\mu_p(\pi)$ is a complex line so that $\mu_p(\pi) \cap \mu_p\left(bB^{n+1}\right)$ is a chain(cf. [Pa3]). Thus the circle

$$\pi \cap bB^{n+1} = \mu_p^{-1}\left(\mu_p(\pi) \cap \mu_p\left(bB^{n+1}\right)\right)$$

is a chain as well. This completes the proof. □

Lemma 12. *Let γ be a chain passing through a point $p \in bB^{n+1}$ and δ_γ be an analytic disk such that*

$$\gamma = \pi \cap bB^{n+1} \quad \text{and} \quad \delta_\gamma = \pi \cap B^{n+1}$$

where π is a complex line. Let θ_γ be the angle between the tangent vector of γ at the point p and a unit vector v_p perpendicular to the complex tangent hyperplane at the point $p \in bB^{n+1}$ and $|\delta_\gamma|$ be the area of the analytic disk δ_γ. Then

$$|a_\gamma| \equiv |\tan\theta_\gamma| \to \infty$$

if and only if

$$|\delta_\gamma| \to 0.$$

Proof. We easily see that

$$\theta_\gamma \to \frac{\pi}{2}$$

if and only if

$$|a_\gamma| \equiv |\tan\theta_\gamma| \to \infty.$$

Note that the complex line π would be on the complex tangent hyperplane of bB^{n+1} at the point p if

$$\theta_\gamma = \frac{\pi}{2}.$$

Since bB^{n+1} is strongly pseudoconvex and B^{n+1} is strongly convex, the complex tangent hyperplane of bB^{n+1} at the point p has no intersection to B^{n+1}. Thus we easily see

$$\theta_\gamma \to \frac{\pi}{2}$$

if and only if

$$|\delta_\gamma| \to 0.$$

This completes the proof. □

We may require the following well-known results in this article(cf. [Kr], [Ra], [Bo]).

Lemma 13 (Lewy, Pinchuk)**.** *Let D, D' be domains with strongly pseudoconvex real analytic boundaries bD, bD' and U be a connected open neighborhood of a point $p \in bD$. Suppose that there is a holomorphic mapping ϕ on $U \cap D$ such that*

$$\phi \in H(U \cap D) \cap C^1\left(U \cap \overline{D}\right), \quad \phi(U \cap bD) \subset bD'$$

and the induced mapping $\phi : U \cap bD \to bD'$ is CR diffeomorphic. Then the mapping ϕ is analytically continued on U, if necessary, shrinking U.

Lemma 14 (Lewy)**.** *Let D be a domain with a strongly pseudoconvex boundary bD and U be an open connected neighborhood of a point $p \in bD$. Suppose that there is a holomorphic mapping ϕ on $U \cap bD$. Then there is an open neighborhood V of the point p such that the mapping ϕ is analytically continued onto*

$$V \cap \overline{D}.$$

Lemma 15 (Wong, Rosay)**.** *Let D be a strongly pseudoconvex bounded domain. Suppose that there is a compact set $K \subset\subset D$ and a sequence $p_j \in K$ and automorphisms $\phi_j \in Aut\,(D)$ such that*

$$\phi_j\,(p_j) \to bD.$$

Then the domain D is biholomorphic to an open unit ball B^{n+1}.

Lemma 16 (Bell-Catlin, Diederich-Fornaess)**.** *Let D, D' be strongly pseudoconvex bounded domains with the boundaries bD, bD' of class C^∞ and $\phi : D \to D'$ be a proper holomorphic mapping. Then $\phi : D \to D'$ is a locally biholomorphic mapping and the induced mapping $\phi : bD \to bD'$ is a locally CR diffeomorphism.*

1. Analytic Continuation on a Sphere

1.1. **Analytic continuation with finiteness.**

Let D be a domain in $\mathbb{C}^{n+1}, n \geq 1$, with real analytic boundary bD. The boundary bD shall be called spherical if, for each point $p \in bD$, there is a connected open neighborhood U of the point p and a biholomorphic mapping ϕ on U such that

$$\phi\,(U \cap bD) \subset bB^{n+1}.$$

Note that a domain D with spherical real analytic boundary is necessarily strongly pseudoconvex.

Lemma 17. *Let p be a point of bB^{n+1} and U be a connected open neighborhood of the point p. Suppose that there is a biholomorphic mapping ϕ on U such that*

$$\phi\left(U \cap bB^{n+1}\right) \subset bB^{n+1}. \tag{4}$$

Then the mapping ϕ is analytically continued on an open neighborhood of the closed ball $\overline{B^{n+1}}$.

Proof. Note that each local automorphism $\varphi \in Aut_p\left(bB^{n+1}\right)$ for any point $p \in bB^{n+1}$ is necessarily birational such that φ is analytically continued on an open neighborhood of $\overline{B^{n+1}}$ as a biholomorphic mapping(cf.[Pa1]).

Let's put $q = \phi\,(p)$. We take a point $r \in bB^{n+1}$ such that $r \neq p$, $r \neq q$. Then we take an automorphism

$$\psi \in Aut_r\left(bB^{n+1}\right)$$

satisfying

$$\psi\,(q) = p$$

so that

$$\varphi \equiv \psi \circ \phi \in Aut_p\left(bB^{n+1}\right).$$

Then the mapping $\psi^{-1}\circ\varphi$ is an automorphism of B^{n+1} and an analytic continuation of the mapping ϕ such that

$$\phi = \psi^{-1}\circ\varphi \quad \text{on} \quad U.$$

This completes the proof. □

Theorem 18. *Let D be a domain in $\mathbb{C}^{n+1}$ with spherical real analytic boundary bD. Suppose that there is a connected open neighborhood U of a point $p\in bD$ and a biholomorphic mapping ϕ on U such that $\phi(U\cap bD)\subset bB^{n+1}$. Then the mapping ϕ is analytically continued along any path on bD as a local biholomorphic mapping.*

Proof. Suppose that the assertion is not true. Then there would be a path $\gamma : [0,1]\to bD$ such that $\gamma(0)\in U\cap bD$ and the germ of a biholomorphic mapping ϕ at the point $\gamma(0)$ is analytically continued along the subpath $\gamma[0,\tau]$ with all $\tau<1$ as a local biholomorphic mapping, but not the whole path $\gamma[0,1]$.

Since bD is spherical, by definition, there exist a connected open neighborhood V of the point $\gamma(1)$ and a biholomorphic mapping φ on V such that

$$\varphi(V\cap bD)\subset bB^{n+1}.$$

We take $\lambda\in[0,1)$ such that

$$\gamma(\tau)\in V\cap bD \quad \text{for all } \tau\in[\lambda,1]$$

and we take a sufficiently small connected open neighborhood W of the point $\gamma(\lambda)$ such that ϕ is analytically continued on $W\subset V$ along the path $\gamma[0,\lambda]$ and

$$\varphi(W)\cap bB^{n+1}\neq\emptyset.$$

Then we have

$$\psi\left(\varphi(W)\cap bB^{n+1}\right)\subset bB^{n+1}$$

where

$$\psi=\phi\circ\varphi^{-1}.$$

By Lemma 17, ψ is analytically continued on an open neighborhood of bB^{n+1} as a local biholomorphic mapping. By abuse of notation, the mapping $\psi\circ\varphi$ is biholomorphic on the open set V such that

$$\psi\circ\varphi=\phi \quad \text{on } W.$$

Thus the germ $\psi\circ\varphi$ at the point $\gamma(1)$ is an analytic continuation of the germ ϕ at the point $\gamma(0)$ along the path $\gamma[0,1]$ as a local biholomorphic mapping. This contradiction completes the proof. □

Lemma 19. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical real analytic boundary bD such that the fundamental group $\pi_1(bD)$ is finite. Suppose that there is a connected open neighborhood U of a point $p\in bD$ and a biholomorphic mapping ϕ on U such that $\phi(U\cap bD)\subset bB^{n+1}$. Then ϕ is analytically continued to a biholomorphic mapping from D onto B^{n+1}.*

Proof. By Lemma 18, the mapping ϕ is analytically continued along any path on bD as a local biholomorphic mapping. Let E be the path space of bD pointed at the point p mod homotopy so that E is a universal covering of bD with a natural CR structure and a CR projection $\varphi : E\to bD$. Then there is a unique CR lift $\psi : E\to bB^{n+1}$ as the analytic continuation of the biholomorphic mapping ϕ. Note that $\psi : E \to bB^{n+1}$ is an open mapping because ϕ and φ are both locally CR diffeomorphisms.

Since bD is finitely connected, E is necessarily compact so that the mapping $\psi : E \to bB^{n+1}$ is surjective. Further, the mapping $\psi : E \to bB^{n+1}$ is a simple covering map because ψ is locally a CR diffeomorphism and the sphere bB^{n+1} is simply connected. Hence there exists a locally biholomorphic mapping $\lambda : bB^{n+1} \to bD$ defined by

$$\lambda = \varphi \circ \psi^{-1} \quad \text{on} \quad bB^{n+1}.$$

By Hartogs extension theorem, the mapping λ uniquely extends to the open ball B^{n+1} as a local biholomorphic mapping and, further, the extension is smooth up to the boundary. Hence λ is well defined as a locally biholomorphic mapping on an open neighborhood of $\overline{B^{n+1}}$ by Lemma 13.

We obtain a proper mapping $\lambda : B^{n+1} \to D$ so that λ is a globally branched covering and the branched locus of λ cannot be bounded by bD. Since $\lambda^{-1} = \phi$ is locally biholomorphic on bD, $\lambda : B^{n+1} \to D$ and $\lambda : bB^{n+1} \to bD$ are finite coverings respectively of D and bD. Since the closed ball $\overline{B^{n+1}}$ has the fixed point property, the mapping $\lambda : B^{n+1} \to D$ is globally one-to-one. Otherwise, there would be a nontrivial deck transform of $\overline{B^{n+1}}$ which is continuous on $\overline{B^{n+1}}$ without a fixed point. Hence the mapping $\lambda^{-1} : D \to B^{n+1}$ is biholomorphic with $\lambda^{-1} = \phi$ on bD. This completes the proof. □

Theorem 20. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with a connected spherical real analytic boundary bD. Suppose that there is a biholomorphic mapping ϕ on a connected open neighborhood U of a point $p \in bD$ satisfying*

$$\phi(U \cap bD) \subset bB^{n+1}$$

such that the analytic continuation of ϕ on the boundary bD yields finitely many germs at each point on bD. Then D is necessarily simply connected and the mapping ϕ is analytically continued to a biholomorphic mapping from D onto B^{n+1}.

Proof. We claim that there is a finite covering space E_1 of bD with a natural CR structure and a CR projection $\varphi_1 : E_1 \to bD$ and a local CR diffeomorphism $\psi_1 : E_1 \to bB^{n+1}$ satisfying the relation $\psi_1 = \phi \circ \varphi_1$. Then the desired result follows from this claim by the same argument in the proof of Lemma 19.

Let E be the path space of bD pointed at the point $p \in bD$ mod homotopy so that E is a universal covering of bD with a natural CR structure and a CR projection $\varphi : E \to bD$. Then there is a unique CR lift $\psi : E \to bB^{n+1}$:

$$\begin{tikzcd} E \arrow[d, "\varphi"'] \arrow[dr, "\psi"] & \\ bD \arrow[r, "\phi"] & bB^{n+1} \end{tikzcd}$$

satisfying the relation $\psi = \phi \circ \varphi$. Note that $\psi : E \to bB^{n+1}$ is an open mapping because ϕ and φ are both local CR diffeomorphisms. Let F be the image of the mapping ψ such that $F = \psi(E)$. Then F is an open subset of bD.

Suppose that $bF \neq \emptyset$. Then we take a point $p \in bF$ and a sequence $p_j \in F$ such that $p_j \to p$. Thus there exists a point $q_j \in bD$ and germs of biholomorphic mappings ϕ_j such that

$$\phi_j(q_j) \to p$$

where ϕ_j are analytic continuations of the mapping ϕ on bD. Since bD is compact, there exist a point $q \in bD$ and a subsequence q_{m_j} of q_j such that $q_{m_j} \to q$. Further, by passing to a subsequence, if necessary, we may assume that

$$\phi^* = \phi_{m_j} \quad \text{for all } m_j$$

because the analytic continuation of the mapping ϕ yields only finitely many germs at the point $q \in bD$. Hence we obtain

$$\phi^* (q) = \lim_{j\to\infty} \phi_{m_j} \left(q_{m_j}\right) = p \in F.$$

This contradiction implies that $bF = \emptyset$, i.e., $\psi(E) = bB^{n+1}$.

For each point $q \in bB^{n+1}$, there is a subset $X_q \subset E$ such that

$$X_q = \{p \in E : \psi(p) = q\}.$$

Since ψ is a CR diffeomorphism, X_q is necessarily a discrete set on E. Then we define a subset $Y_q \subset bD$ such that

$$Y_q = \{\varphi(p) \in bD : p \in X_q\}.$$

Suppose that Y_q has an accumulation point $y \in bD$. Then there is a sequence of point $p_j \in bD$ satisfying

$$p_j \to y \quad \text{and} \quad p_j \neq y,$$

and biholomorphic mappings ϕ_j such that

$$\phi_j (p_j) = q$$

where the mapping ϕ at the point p_j is the analytic continuation of the mapping ϕ. Because the analytic continuation of ϕ yields only finitely many germs at the point y, we can take a subsequence of ϕ and a biholomorphic mapping ϕ^* such that $\phi^* = \phi_{m_j}$. Then we have

$$\phi^* (y) = \lim_{j\to\infty} \phi_{m_j} \left(p_{m_j}\right) = q.$$

Since ϕ^* is locally biholomorphic, it is impossible that $\phi^*\left(p_{m_j}\right) = q = \phi^*(y)$ and $p_{m_j} \to y, p_{m_j} \neq y$ at the same time. Thus we find that the set Y_q is finite.

Therefore, the analytic continuation of the mapping ϕ on bD is mapped to a point of bB^{n+1} only at finitely many points of bD. Since the analytic continuation of the mapping ϕ yields finitely many germs at each point on bD, only finitely many germs of the analytic continuation of the mapping ϕ on bD are mapped to each point of bB^{n+1}. Then, by the compactness of bB^{n+1}, we obtain a finite covering space E_1 of bD satisfying all conditions in the claim. This completes the proof. ☐

1.2. **First Dogginal Lemma.**

Lemma 21 (First Scaling Lemma)**.** *Let p be a point of the boundary bB^{n+1} and p_j, $j \in \mathbb{N}^+$, be a sequence of points of bB^{n+1} such that $p_j \neq p$ for all j and $p_j \to p$ as $j \to \infty$ to a direction transversal to the complex tangent hyperplane at the point $p \in bB^{n+1}$. Let ε_j be the euclidean distance between the two points p_j and p, and δ_j be the analytic disk*

$$\delta_j = \pi_j \cap B^{n+1}$$

where π_j is the complex line passing through the two points p_j and p. Suppose that there is a sequence p'_j of points of bB^{n+1} satisfying

$$p'_j \to p' \in bB^{n+1},$$

and a sequence of biholomorphic automorphisms $\phi_j \in Aut\left(B^{n+1}\right)$ satisfying

$$\phi_j\left(p'_j\right) = p_j$$

such that the sequence ϕ_j converges to a constant mapping and the area $\left|\phi_j^{-1}\left(\delta_j\right)\right|$ of the analytic disks

$$\phi_j^{-1}\left(\delta_j\right)$$

is bounded from the below, i.e., there is a real number $c > 0$ satisfying

$$\left|\phi_j^{-1}\left(\delta_j\right)\right| \geq c.$$

Then there is a subsequence ϕ_{m_j} and a sequence of local automorphisms $\sigma_j \in Aut_{p_{m_j}}\left(B^{n+1}\right)$ such that

$$U_{\sigma_j} = id_{n\times n}, \quad \rho_{\sigma_j} = \varepsilon_{m_j}, \quad a_{\sigma_j} = 0, \quad r_{\sigma_j} = 0$$

and the composition

$$\sigma_j^{-1} \circ \phi_{m_j} : B^{n+1} \to B^{n+1}$$

uniformly converges to an automorphism of the unit ball B^{n+1}.

Proof. Note that there is a subsequence ϕ_{m_j} which converges to the point $p \in bB^{n+1}$ uniformly on every compact subset of the unit ball B^{n+1}. We take a point $p' \in bB^{n+1}$ such that $p' \neq p$ and

$$p' \notin \overline{\left\{p_{m_j} : j \in \mathbb{N}^+\right\}},$$

if necessary, passing to a subsequence.. Then, by Lemma 8, there is a unique decomposition of the automorphism ϕ_{m_j} such that

$$\phi_{m_j} = \varphi_j \circ \psi_j$$

where

$$\varphi_j \in Aut_{p_{m_j}}\left(bB^{n+1}\right), \quad \psi_j \in Aut_{p'}\left(bB^{n+1}\right)$$

and the local automorphism ψ_j acts trivially on the complex tangent hyperplane of bB^{n+1} at the fixed point p'.

Let $U_{\psi_j}, \rho_{\psi_j}, a_{\psi_j}, r_{\psi_j}$ be the normalizing parameters of the local automorphism ψ_j. Since the local automorphism ψ_j acts trivially on the complex tangent hyperplane of bB^{n+1} at the fixed point p', we obtain

$$U_{\psi_j} = id_{n\times n}, \quad \rho_{\psi_j} = 1.$$

Since the sequence ϕ_{m_j} uniformly converges to the point p, there is a real number $e > 0$ such that

$$\left|a_{\psi_j}\right| \leq e, \quad \left|r_{\psi_j}\right| \leq e. \tag{5}$$

Let π_j be the complex line passing through the two points p and p_{m_j}. Since the points p_j converges to the point p to a direction transversal to the complex tangent hyperplane at the point $p \in bB^{n+1}$, the analytic disks

$$\delta_j = \pi_j \cap B^{n+1}$$

is uniformly bounded in their area from the below by a positive number.

Let $U_{\varphi_j}, \rho_{\varphi_j}, a_{\varphi_j}, r_{\varphi_j}$ be the normalizing parameters of the local automorphism φ_j. Note that the analytic disk

$$\phi_{m_j}^{-1}(\delta_j)$$

is mapped by ϕ_{m_j} onto the analytic disk δ_j, where the areas of the analytic disks in both classes

$$\left|\phi_{m_j}^{-1}(\delta_j)\right| \quad \text{and} \quad |\delta_j|$$

are bounded from the below. Since the chain $\phi_{m_j}^{-1}(b\delta_j)$ is mapped by ϕ_{m_j} to the chain $b\delta_j$, there is a real number $e > 0$, if necessary, increasing e, such that

$$\left|a_{\varphi_j}\right| \le e. \tag{6}$$

Since the area of the analytic disks δ_j is bounded by a positive number, the point p is attracted to the center of the local automorphism $Aut_{p_{m_j}}\left(bB^{n+1}\right)$. Then, by passing to a subsequence, if necessary and increasing e, there is a real number $e > 0$ such that

$$\left|r_{\varphi_j}\right| \le e \tag{7}$$

and

$$\rho_{\varphi_j} \to 0 \quad \text{as } j \to \infty.$$

Since bB^{n+1} is strongly pseudoconvex, we obtain

$$\left|U_{\varphi_j}\right| = 1.$$

Let η_j be a local automorphism in $Aut_{p_{m_j}}\left(bB^{n+1}\right)$ defined by the normalizing parameters

$$U_{\eta_j} = id_{n\times n}, \quad \rho_{\eta_j} = \rho_{\varphi_j}, \quad a_{\eta_j} = 0, \quad r_{\eta_j} = 0.$$

Then the composition

$$\varphi_j' \equiv \eta_j^{-1} \circ \varphi_j \in Aut_{p_{m_j}}\left(bB^{n+1}\right)$$

has the same normalizing parameters of the mapping φ_j except for $\rho_{\varphi_j'} = 1$, i.e.,

$$U_{\varphi_j'} = U_{\varphi_j}, \quad \rho_{\varphi_j'} = 1, \quad a_{\varphi_j'} = a_{\varphi_j}, \quad r_{\varphi_j'} = r_{\varphi_j}.$$

Therefore, by passing to a subsequence, if necessary, the sequence

$$\tau_j \equiv \eta_j^{-1} \circ \phi_{m_j} : B^{n+1} \to B^{n+1}$$

converges by Hurwitz theorem to a locally biholomorphic mapping. Since τ_j are automorphism of B^{n+1}, the sequence τ_j converges to an automorphism of B^{n+1}.

Let $q \in \overline{B^{n+1}}$ be the limit point of the sequence

$$\tau_j\left(p_{m_j}'\right) \to q \in \overline{B^{n+1}}.$$

We set

$$\lambda_j = \frac{\varepsilon_{m_j}}{\rho_{\varphi_j}}.$$

The sequence λ_j converges to the distance between the two distinct points q and $p \in bB^{n+1}$ so that there is a real number $e > 0$ such that

$$e^{-1} \le |\lambda_j| \le e, \tag{8}$$

if necessary, increasing e.

Let σ_j be a local automorphism in $Aut_{p_{m_j}}\left(bB^{n+1}\right)$ defined by the normalizing parameters

$$U_{\sigma_j} = id_{n\times n}, \quad \rho_{\sigma_j} = \varepsilon_{m_j}, \quad a_{\sigma_j} = 0, \quad r_{\sigma_j} = 0.$$

Then the composition

$$\varphi_j'' \equiv \sigma_j^{-1} \circ \varphi_j \in Aut_{p_{m_j}}\left(bB^{n+1}\right)$$

has the same normalizing parameters of the mapping φ_j except for $\rho_{\varphi_j''} = \lambda_j^{-1}$, i.e.,

$$U_{\varphi_j''} = U_{\varphi_j}, \quad \rho_{\varphi_j''} = \lambda_j^{-1}, \quad a_{\varphi_j''} = a_{\varphi_j}, \quad r_{\varphi_j''} = r_{\varphi_j}.$$

Therefore, by the estimates 5, 6, 7, 8, the sequence

$$\tau_j' \equiv \sigma_j^{-1} \circ \phi_{m_j} : B^{n+1} \to B^{n+1}$$

uniformly converges to an automorphism of B^{n+1}. This completes the proof. □

Lemma 22. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical real analytic boundary bD. Suppose that there is a connected open neighborhood U of a point $p \in bD$ and a biholomorphic mapping ϕ on U such that $\phi(U \cap bD) \subset bB^{n+1}$ and the inverse mapping ϕ^{-1} on bB^{n+1} is analytically continued along every chain of bB^{n+1}. Then the mapping ϕ is analytically continued to a biholomorphic mapping from D onto B^{n+1}.*

Proof. The chain on bB^{n+1} is characterized to be the intersection of a complex line on bB^{n+1}. Thus the chains on bB^{n+1} form a continuous family so that the analytic continuity of the inverse mapping ϕ^{-1} along every chain on bB^{n+1} is equivalent to the analytic continuity along any path on bB^{n+1}.

Note that the inverse mapping ϕ^{-1} is analytically continued along any path on bB^{n+1} and, by Hartogs extension theorem, the branching locus of a proper mapping cannot be bounded by bD. Since bB^{n+1} is simply connected, by the monodromy theorem, the mapping ϕ^{-1} is analytically continued to, by abuse of notation, a locally biholomorphic proper mapping $\phi^{-1} : B^{n+1} \to D$ such that $\phi^{-1}\left(bB^{n+1}\right) = bD$. By the fixed point property of the closed ball $\overline{B^{n+1}}$, the proper mapping $\phi^{-1} : B^{n+1} \to D$ is globally one-to-one so that the mapping $\phi^{-1} : B^{n+1} \to D$ is biholomorphic. This completes the proof. □

Lemma 23. *Let D be a domain in $\mathbb{C}^{n+1}$ with spherical real analytic boundary bD and U be a connected open neighborhood of a point $p \in bD$. For a chain γ on bD passing through the point p and tangential to the direction with an angle θ_γ with respect to a unit vector v_p at the point p perpendicular to the complex tangent hyperplane, we denote $|a_\gamma| \equiv |\tan\theta_\gamma|$. Then there is a real number $e > 0$ such that every chain γ passing through the point p with $|a_\gamma| \geq e$ is the boundary of a nonsingular analytic disk $\delta_\gamma \subset U \cap D$ such that*

$$\gamma = \overline{\delta_\gamma} \cap bD.$$

Proof. We take a biholomorphic mapping ϕ on the open set U, if necessary, shrinking U such that

$$\phi(U \cap bD) \subset bB^{n+1}.$$

Every chain λ on bB^{n+1} is an intersection with a complex line π_λ such that

$$\lambda = \pi_\lambda \cap bB^{n+1}.$$

For a sufficiently large real number $e > 0$, each chain γ with $|a_\gamma| \geq e$ is obtain by the relation

$$\gamma = \phi^{-1}(\lambda)$$

where the chain λ on bB^{n+1} satisfies the condition

$$\pi_\lambda \cap B^{n+1} \subset \phi(U \cap D).$$

Then we take

$$\delta_\gamma = \phi^{-1}\left(\pi_\lambda \cap B^{n+1}\right).$$

This completes the proof. ☐

Lemma 24 (First Dogginal Lemma)**.** *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical real analytic boundary bD and ϕ be a biholomorphic mapping on a connected open neighborhood U of a point $p \in bB^{n+1}$ satisfying*

$$\phi\left(U \cap bB^{n+1}\right) \subset bD.$$

Suppose that there is a chain-segment $\gamma : [0,1] \to bB^{n+1}$ such that $\gamma(0) \in U \cap bB^{n+1}$ and the mapping ϕ is analytically continued along the subpath $\gamma[0,\tau]$ for all $\tau < 1$, but not the whole path $\gamma[0,1]$ as a local biholomorphic mapping. Let π be the complex line containing the chain-segment $\gamma[0,1]$. Then there is an open neighborhood V along the path $\gamma[0,1]$ such that

1. $\gamma[0,\tau] \subset V$ *for all* $\tau < 1$,
2. $bV \cap \pi \cap B(\gamma(1);\delta)$ *is an angle for a sufficient small* $\delta > 0$, *which contains the chain-segment* $\gamma[0,1]$,
3. $bV \cap bB^{n+1} \cap B(\gamma(1);\delta)$ *is slanted paraboloid for a sufficiently small* $\delta > 0$, *which smoothly touches the complex tangent hyperplane at the point* $\gamma(1)$,
4. *the mapping ϕ is analytically continued on V as a local biholomorphic mapping.*

Proof. By the analytic continuation of the mapping ϕ along the subpath $\gamma[0,\tau]$ for all $\tau < 1$, there is a path $\phi \circ \gamma : [0,1) \to bD$. Then we consider the following sequences

$$\begin{aligned} p_j &= \gamma\left(1 - \frac{1}{j}\right), \quad \text{for } j \in \mathbb{N}^+, \\ p'_j &= \phi \circ \gamma\left(1 - \frac{1}{j}\right), \quad \text{for } j \in \mathbb{N}^+. \end{aligned}$$

Since bD is compact, there is a subsequence p'_{m_j} and a point $p' \in bD$ such that

$$p'_{m_j} \to p'.$$

By Theorem 18, the mapping ϕ^{-1} is analytically continued along the path $\phi \circ \gamma[0,1) \subset bD$.

Let φ_j be the analytic continuation of the mapping ϕ^{-1} at the point p'_{m_j} along the path $\phi \circ \gamma[0,1) \subset bD$. By Theorem 18, there is an open neighborhood W of the point p' such that φ_j is locally biholomorphic on an open neighborhood of $W \cap bD$.

By Lemma 14, we may assume that φ_j is holomorphic on $W \cap D$, if necessary, shrinking W.

Since bD is spherical, there is an open neighborhood W of the point p', if necessary, shrinking W, and a biholomorphic mapping φ on W such that

$$\varphi\left(W \cap bD\right) \subset bB^{n+1}.$$

Then, by Lemma 17, the compositions

$$\phi_j \equiv \varphi_j \circ \varphi^{-1} : \varphi\left(W\right) \cap bB^{n+1} \subset bB^{n+1}$$

are analytically continued, by abuse of notation, to automorphisms ϕ_j of the unit ball B^{n+1}. Without loss of generality, we may assume that the sequence ϕ_j converges to a holomorphic mapping uniformly on every compact subset of B^{n+1}.

We set

$$p''_{m_j} \equiv \varphi\left(p'_{m_j}\right) \in bB^{n+1} \quad \text{and} \quad p'' \equiv \varphi\left(p'\right)$$

so that

$$p''_{m_j} \to p''.$$

Hence the relation

$$\phi_j\left(p''_{m_j}\right) = p_{m_j}$$

yields

$$\phi_j\left(p''_{m_j}\right) \to p.$$

By Lemma 9, the mapping ϕ_j converges to either a constant mapping or an automorphism of B^{n+1}. We claim that the sequence ϕ_j converges to the point p uniformly on every compact subset of B^{n+1}. Otherwise, the sequence ϕ_j converges to an automorphism of B^{n+1} so that the sequence

$$\varphi_j = \phi_j \circ \varphi$$

converges to a biholomorphic mapping on an open neighborhood W of the point p'. Then there is a real number δ such that the mapping φ_j and its inverse φ_j^{-1} are analytically continued respectively on

$$B\left(p'_{m_j}; \delta\right) \quad \text{and} \quad B\left(p_{m_j}; \delta\right)$$

as a locally biholomorphic mapping. Thus the mapping $\phi = \varphi_j^{-1}$ is biholomorphic on $B\left(p_{m_j}; \delta\right)$ for every point $p_{m_j} \to p$. This is impossible by the hypothesis on the mapping ϕ. Thus, by Lemma 9, the sequence ϕ_j converges to the point p on every compact subset of B^{n+1}.

Let π_j be the complex line passing through the two points p and p_{m_j}, and δ_j be the analytic disk

$$\delta_j = \pi_j \cap B^{n+1}.$$

Let ε_j be the euclidean length between the two points p and p_{m_j}. Note that the area $\left|\phi_j^{-1}\left(\delta_j\right)\right|$ of the analytic disk

$$\phi_j^{-1}\left(\delta_j\right)$$

is bounded from the below. Otherwise, by Lemma 23, the mapping $\phi = \varphi_j^{-1}$ is analytically continued over the point p as a locally biholomorphic mapping. Therefore, by First Scaling Lemma, there is a sequence $\sigma_j \in Aut_{p_{m_j}}\left(bB^{n+1}\right)$ such that

$$U_{\sigma_j} = id_{n\times n}, \quad \rho_{\sigma_j} = \varepsilon_j, \quad a_{\sigma_j} = 0, \quad r_{\sigma_j} = 0$$

and the sequence

$$\psi_j \equiv \sigma_j^{-1} \circ \phi_j$$

converges to an automorphism of B^{n+1}.

Note that there is a real number $\delta > 0$ such that the mapping φ and its inverse φ^{-1} are biholomorphically continued respectively on

$$B\left(p'_{m_j};\delta\right) \quad \text{and} \quad B\left(p''_{m_j};\delta\right),$$

if necessary, passing to a subsequence. Further, there is a real number $\delta > 0$ such that the mapping ψ_j and its inverse ψ_j^{-1} are biholomorphically continued respectively to

$$B\left(p'_{m_j};\delta\right) \quad \text{and} \quad B\left(p_{m_j};\delta\right),$$

if necessary, shrinking δ. Then the mapping

$$\begin{aligned} \phi &= \varphi_j^{-1} \\ &= \varphi^{-1} \circ \psi_j^{-1} \circ \sigma_j^{-1} \end{aligned}$$

is biholomorphically continued on the open neighborhood

$$\sigma_j\left(B\left(p_{m_j};\delta\right)\right).$$

For a canonical normalizing mapping $\mu_{p_{m_j}}$, we obtain

$$\sigma'_j \equiv \mu_{p_{m_j}} \circ \sigma_j \circ \mu_{p_{m_j}}^{-1} : \begin{cases} z^* = \sqrt{\varepsilon_j} z \\ w^* = \varepsilon_j w \end{cases} .$$

Since $p_{m_j} \to p$, by Lemma 5, there is a real number $\delta > 0$ such that the mapping $\mu_{p_{m_j}}$ and its inverse $\mu_{p_{m_j}}^{-1}$ are biholomorphically continued respectively to

$$B\left(p_{m_j};\delta\right) \quad \text{and} \quad B\left(0;\delta\right).$$

Hence the mapping

$$\phi \circ \mu_{p_{m_j}}^{-1} = \varphi^{-1} \circ \psi_j^{-1} \circ \mu_{p_{m_j}}^{-1} \circ \sigma_j'^{-1}$$

is biholomorphically continued on

$$\sigma'_j\left(B\left(0;\delta\right)\right).$$

Since $p_{m_j} \to p$, by Lemma 5, the canonical normalizing mapping $\mu_{p_{m_j}}$ uniformly converges to the mapping μ_p so that the mapping ϕ is biholomorphically continued near the point p_{m_j} on

$$\mu_{p_{m_j}}^{-1} \circ \sigma'_j\left(B\left(0;\delta\right)\right).$$

Therefore, the analytically continued region of the mapping ϕ along the chain $\gamma[0,1) \subset bB^{n+1}$ contains an open set along the chain $\gamma[0,1]$ which touches to the point $\gamma(1)$ by an edge shape transversal to bB^{n+1} and by a slanted paraboloid shape on bB^{n+1}. This completes the proof. $\square$

1.3. **Doggaebi variety on a sphere.** Let ϕ be a biholomorphic mapping on an open neighborhood U of a point $p \in bD$ satisfying $\phi(U \cap bD) \subset bB^{n+1}$. Let $L \subset bB^{n+1}$ be the singular locus of the analytic continuation of the mapping ϕ^{-1}, which shall be called the Doggaebi variety associated to the mapping ϕ.

Lemma 25 (Second Scaling Lemma)**.** *Let* p *be a point of the boundary* bB^{n+1} *and* p_j, $j \in \mathbb{N}^+$, *be a sequence of points of* bB^{n+1} *such that* $p_j \neq p$ *for all* j *and* $p_j \to p$ *as* $j \to \infty$ *to a direction tangential to the complex tangent hyperplane at the point* $p \in bB^{n+1}$. *Let* ε_j *be the euclidean distance between the two points* p_j *and* p, *and* δ_j *be the analytic disk*

$$\delta_j = \pi_j \cap B^{n+1}$$

where π_j *is the complex line passing through the two points* p_j *and* p. *Suppose that there is a sequence* p'_j *of points of* bB^{n+1} *satisfying*

$$p'_j \to p' \in bB^{n+1},$$

and a sequence of biholomorphic automorphisms $\phi_j \in Aut\left(B^{n+1}\right)$ *satisfying*

$$\phi_j\left(p'_j\right) = p_j$$

such that the area $\left|\phi_j^{-1}(\delta_j)\right|$ *of the analytic disks*

$$\phi_j^{-1}(\delta_j)$$

is bounded from the below, i.e., there is a real number $c > 0$ *satisfying*

$$\left|\phi_j^{-1}(\delta_j)\right| \geq c.$$

Then there is a subsequence ϕ_{m_j} *and a sequence of local automorphisms* $\sigma_j \in Aut_{p_{m_j}}\left(bB^{n+1}\right)$ *such that*

$$U_{\sigma_j} = id_{n\times n}, \quad \rho_{\sigma_j} = \varepsilon_{m_j}^2, \quad a_{\sigma_j} = 0, \quad r_{\sigma_j} = 0$$

and the composition

$$\sigma_j^{-1} \circ \phi_{m_j} : B^{n+1} \to B^{n+1}$$

uniformly converges to an automorphism of the unit ball B^{n+1}.

Proof. Note that there is a subsequence ϕ_{m_j} which converges to the point $p \in bB^{n+1}$ uniformly on an open neighborhood of the closed ball $\overline{B^{n+1}}$. We take a point $p'' \in bB^{n+1}$ such that $p'' \neq p$ and

$$p'' \notin \overline{\left\{p_{m_j} : j \in \mathbb{N}^+\right\}},$$

if necessary, passing to a subsequence.. Then, by Corollary 8, there is a unique decomposition of the automorphism ϕ_{m_j} such that

$$\phi_{m_j} = \varphi_j \circ \psi_j$$

where

$$\varphi_j \in Aut_{p_{m_j}}\left(bB^{n+1}\right), \quad \psi_j \in Aut_{p''}\left(bB^{n+1}\right)$$

and the local automorphism ψ_j acts trivially on the tangent hyperplane of bB^{n+1} at the fixed point p'.

Let $U_{\psi_j}, \rho_{\psi_j}, a_{\psi_j}, r_{\psi_j}$ be the normalizing parameters of the local automorphism ψ_j. Since the local automorphism ψ_j acts trivially on the tangent hyperplane of bB^{n+1} at the fixed point p'', we obtain

$$U_{\psi_j} = id_{n\times n}, \quad \rho_{\psi_j} = 1.$$

Since the sequence ϕ_{m_j} uniformly converges to the point p, there is a real number $e > 0$ such that

$$\left|a_{\psi_j}\right| \leq e, \quad \left|r_{\psi_j}\right| \leq e. \tag{9}$$

Let π_j be the complex line passing through the two points p and p_{m_j}, and δ_j be the analytic disks

$$\delta_j = \pi_j \cap B^{n+1}.$$

Let σ_j be a local automorphism in $Aut_{p_{m_j}}\left(bB^{n+1}\right)$ defined by the normalizing parameters

$$U_{\sigma_j} = id_{n\times n}, \quad \rho_{\sigma_j} = \varepsilon_{m_j}^2, \quad a_{\sigma_j} = 0, \quad r_{\sigma_j} = 0.$$

Since the points p_j converges to the point p to a direction tangential to the complex tangent hyperplane at the point $p \in bB^{n+1}$, the area $\left|\sigma_j^{-1}\left(\delta_j\right)\right|$ of the analytic disk

$$\sigma_j^{-1}\left(\delta_j\right)$$

is bounded from below.

Let $U_{\varphi_j}, \rho_{\varphi_j}, a_{\varphi_j}, r_{\varphi_j}$ be the normalizing parameters of the local automorphism φ_j. Note that the analytic disk

$$\phi_{m_j}^{-1}\left(\delta_j\right)$$

is mapped by $\sigma_j^{-1} \circ \phi_{m_j}$ onto the analytic disk

$$\sigma_j^{-1}\left(\delta_j\right)$$

where the areas of the analytic disks in both classes are bounded from below. Further, the normalizing parameters $a_{\varphi'_j}, r_{\varphi'_j}$ of the composition $\varphi'_j = \sigma_j^{-1} \circ \varphi_j$ is the same value of $a_{\varphi_j}, r_{\varphi_j}$, i.e.,

$$a_{\varphi'_j} = a_{\varphi_j}, \quad r_{\varphi'_j} = r_{\varphi_j}$$

so that there is a real number $e > 0$, if necessary, increasing e, such that

$$\left|a_{\varphi_j}\right| \leq e. \tag{10}$$

Since the area of the analytic disks $\sigma_j^{-1}\left(\delta_j\right)$ is bounded by a positive number, the point p should be attracted to the center of the local automorphism $Aut_{p_{m_j}}\left(bB^{n+1}\right)$. Hence, by passing to a subsequence, if necessary and increasing e, there is a real number $e > 0$ such that

$$\left|r_{\varphi_j}\right| \leq e \tag{11}$$

and

$$\rho_{\varphi_j} \to 0 \quad \text{as } j \to \infty.$$

Since bB^{n+1} is strongly pseudoconvex, we obtain

$$\left|U_{\varphi_j}\right| = 1.$$

We set

$$\lambda_j = \frac{\varepsilon_{m_j}^2}{\rho_{\varphi_j}}$$

so that there is a real number $e > 0$ such that

$$e^{-1} \leq |\lambda_j| \leq e, \tag{12}$$

if necessary, increasing e. Then the composition

$$\varphi_j' \equiv \sigma_j^{-1} \circ \varphi_j \in Aut_{p_{m_j}}\left(bB^{n+1}\right)$$

has the same normalizing parameters of the mapping φ_j except for $\rho_{\varphi_j'} = \lambda_j^{-1}$, i.e.,

$$U_{\varphi_j'} = U_{\varphi_j}, \quad \rho_{\varphi_j'} = \lambda_j^{-1}, \quad a_{\varphi_j'} = a_{\varphi_j}, \quad r_{\varphi_j'} = r_{\varphi_j}.$$

Therefore, by the estimates 9, 10, 11, 12, the sequence

$$\tau_j \equiv \sigma_j^{-1} \circ \phi_{m_j} : B^{n+1} \to B^{n+1}$$

uniformly converges to an automorphism of B^{n+1}. This completes the proof. □

Theorem 26. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical real analytic boundary bD. Suppose that there is a biholomorphic mapping ϕ on a connected open neighborhood U of a point $p \in bD$ satisfying*

$$\phi(U \cap bD) \subset bB^{n+1}.$$

Then the Doggaebi variety L associated to the mapping ϕ is a finite subset of bB^{n+1} such that the inverse mapping ϕ^{-1} is analytically continued along any piecewise chain curve on $bB^{n+1} \backslash L$ as a locally biholomorphic mapping.

Proof. By First Dogginal Lemma, the singular locus of the analytic continuation of the inverse mapping ϕ^{-1} on bB^{n+1} is an integral manifold of the subdistribution of the complex tangent hyperplanes on bB^{n+1} in its smooth part. Thus the mapping ϕ^{-1} cannot be branched on bB^{n+1} by a branching locus passing through the boundary bB^{n+1}. Otherwise, the intersection of the nontrivial branch of the mapping ϕ^{-1} to the boundary bB^{n+1} would be a nontrivial complex submanifold on bB^{n+1}. Hence the singular locus of the analytic continuation of the mapping ϕ^{-1} is well defined so that the inverse mapping ϕ^{-1} is analytically continued on $bB^{n+1} \backslash L$ as a locally biholomorphic mapping.

We take a chain-segment $\gamma : [0,1] \to bB^{n+1}$ with $\gamma(1) = p \in L$ such that a germ of the mapping ϕ^{-1} at the point $\gamma(0)$ is analytically continued along all subpath $\gamma[0,\tau]$ with $\tau < 1$, but not the whole path $\gamma[0,1]$. Note that the distribution of the complex tangent hyperplanes on bB^{n+1} is maximally nonintegrable. Thus, by First Dogginal Lemma, the singular locus L of the analytic continuation of the mapping ϕ^{-1} cannot bound the open region of the analytic continuation of the mapping ϕ^{-1}. Then the mapping ϕ^{-1} is analytically continued on the opposite side of the complex tangent hyperplane H_p at the point $p \in bB^{n+1}$. Thus there is a complex line π passing through the point p and an open neighborhood U of the point p such that

$$\pi \cap bB^{n+1}$$

is a chain on bB^{n+1} satisfying

$$L \cap U \cap \pi \cap bB^{n+1} = \{p\}.$$

We claim that, if necessary, shrinking U,

$$L\cap U\cap bB^{n+1}=\{p\}$$

so that the singular locus L is a finite set on bB^{n+1}. By First Dogginal Lemma, there is a sequence $p_j\in bB^{n+1}\backslash L$ such that $p_j\to p$ and the sequence p_j converges to the point p to a direction tangential to the complex tangent hyperplane at the point $p\in L\subset bB^{n+1}$. Since bD is compact, there is a subsequence p_{m_j} and a point $p'\in bD$ such that

$$p'_{m_j}\equiv\phi^{-1}\left(p_{m_j}\right)\to p'.$$

Let ϕ_j be the germ of the mapping ϕ at the point $p'_{m_j}\in bD$ such that

$$\phi_j\left(p'_{m_j}\right)=p_{m_j}.$$

Since bD is spherical, there is an open neighborhood W of the point p' and a biholomorphic mapping ψ on W such that

$$\psi\left(W\cap bD\right)\subset bB^{n+1}.$$

Then the compositions $\varphi_j\equiv\phi_j\circ\psi^{-1}$ satisfy

$$\varphi_j\left(\psi\left(W\right)\cap bB^{n+1}\right)\subset bB^{n+1},$$

if necessary, shrinking W so that, by abuse of notation, the mapping φ_j is an automorphism of the unit ball B^{n+1}.

Let π_j be the complex line passing through the points p_{m_j} and p, and δ_j be the analytic disk

$$\delta_j=\pi_j\cap B^{n+1}.$$

Note that the area $\left|\varphi_j^{-1}\left(\delta_j\right)\right|$ of the analytic disk $\varphi_j^{-1}\left(\delta_j\right)$ is bounded from the below. Otherwise, $p\in bB^{n+1}\backslash L$. Further, the sequence φ_j converges to the point p uniformly on every compact subset of B^{n+1}, if necessary, passing to a subsequence. Otherwise, the sequence φ_j converges to an automorphism of the unit ball B^{n+1} so that $p\in bB^{n+1}\backslash L$. Let ε_j be the euclidean length between the two points p and p_{m_j}. Then, by Second Scaling Lemma, there is a sequence of local automorphisms $\sigma_j\in Aut_{p_{m_j}}\left(bB^{n+1}\right)$ such that

$$U_{\sigma_j}=id_{n\times n},\quad \rho_{\sigma_j}=\varepsilon_j^2,\quad a_{\sigma_j}=0,\quad r_{\sigma_j}=0$$

and the composition

$$\tau_j\equiv\sigma_j^{-1}\circ\varphi_j:B^{n+1}\to B^{n+1}$$

uniformly converges to an automorphism of the unit ball B^{n+1}. Thus there is a positive real number $\delta>0$ such that the mapping τ_j and its inverse τ_j^{-1} are analytically continued respectively on

$$B\left(p''_{m_j};\delta\right)\quad\text{and}\quad B\left(p_{m_j};\delta\right)$$

where

$$p''_{m_j}=\psi\left(p'_{m_j}\right).$$

Therefore the mapping ϕ^{-1} at the point p_{m_j} :

$$\begin{aligned} \phi^{-1} &= \phi_j^{-1} \\ &= \psi^{-1} \circ \tau_j^{-1} \circ \sigma_j^{-1} \end{aligned}$$

is analytically continued to the open neighborhood

$$\sigma_j \left(B \left(p_{m_j}; \delta \right) \right).$$

By the canonical normalizing mapping $\mu_{p_{m_j}}$, we obtain

$$\sigma_j' \equiv \mu_{p_{m_j}} \circ \sigma_j \circ \mu_{p_{m_j}}^{-1} : \begin{cases} z^* = \varepsilon_j z \\ w^* = \varepsilon_j^2 w \end{cases}.$$

Hence the mapping ϕ^{-1} at the point p_{m_j} is analytically continued to the open neighborhood

$$\mu_{p_{m_j}}^{-1} \circ \sigma_j' \left(B \left(0; \delta \right) \right),$$

if necessary, shrinking δ. Since ε_j is the euclidean length between the two points p and p_{m_j}, the mapping ϕ^{-1} is analytically continued on an open region which touches to the point p to a converging direction to the sequence p_j by an edge shape tangential to the complex tangent hyperplane H_p at the point p and a $\sqrt{|x|}$ curve shape normal to H_p on bB^{n+1}.

Therefore, the singular locus L of the analytic continuation of the mapping ϕ^{-1} is isolated to the direction of the complex tangent hyperplane at each point of L. Since the boundary bB^{n+1} is compact, the singular locus L is a finite subset of bB^{n+1}. This completes the proof. □

Lemma 27. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical real analytic boundary bD. Then every chain on bD is extended each direction infinitely in its euclidean length.*

Proof. Suppose that the assertion is not true. Then there would be a path $\gamma : [0,1] \to bD$ such that the subpath $\gamma[0,\tau]$ is a chain-segment for all $\tau < 1$ but the whole path $\gamma[0,1]$ is not a chain segment.

Since bD is spherical, by definition, there exist a connected open neighborhood U of the point $\gamma(1)$ and a biholomorphic mapping ϕ on U such that $\phi(U \cap bD) \subset bB^{n+1}$. There is a unique closed circle χ on bB^{n+1} such that χ is a chain on bB^{n+1} and $\phi \circ \gamma[0,\tau] \subset \chi$ for all $\tau < 1$(cf. [Pa3]).

Then the inverse image $\phi^{-1}(\chi \cap \phi(U))$ under the biholomorphic mapping ϕ is a chain on bB^{n+1}(cf. [Pa3]) such that

$$\gamma[0,1] \cap U \subset \phi^{-1}(\chi \cap \phi(U)).$$

This is a contradiction. This completes the proof. □

Theorem 28. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical real analytic boundary bD. Suppose that there is a biholomorphic mapping ϕ on a connected open neighborhood U of a point $p \in bD$ satisfying*

$$\phi(U \cap bD) \subset bB^{n+1}.$$

Let E be the path space of bD pointed at the point $p \in bD$ mod homotopic relation so that E be a universal covering of bD with a natural CR structure and a natural

CR covering map $\psi : E \to bD$. *Then there is a unique CR equivalence* $\varphi : E \to bB^{n+1} \backslash L$ *commuting the diagram*

$$\begin{tikzcd}
E \arrow[d, "\psi"'] \arrow[dr, "\varphi"] & \\
bD \arrow[r, "\phi"] & bB^{n+1} \backslash L
\end{tikzcd}$$

where $L \subset bB^{n+1}$ *is the Doggaebi variety associated to the mapping* ϕ.

Proof. We obtain the set $bB^{n+1} \backslash L$ when the mapping ϕ is analytically continued along chains on bD. Since the mapping $\varphi : E \to bB^{n+1}$ is naturally induced by the analytic continuation of the mapping ϕ on bD(cf. Theorem 18), we obtain

$$bB^{n+1} \backslash L \subset \varphi(E)$$

Let $\gamma : [0, 1] \to bD$ be a path such that $\gamma(0) = p$. Then, for a sufficiently small $\varepsilon > 0$, there is an ε open neighborhood V of the path $\gamma[0, 1]$ such that the mapping ϕ is analytically continued on V as a local biholomorphic mapping. Then we take a piecewise chain curve $\eta : [0, 1] \to bD$(cf. [Pa3]) such that

$$\eta(0) = \gamma(0), \quad \eta(1) = \gamma(1)$$

and

$$\eta[0, 1] \subset V.$$

Further, we may take a continuous function $\Gamma : [0, 1] \times [0, 1] \to V$ such that

$$\Gamma(0, \tau) = \gamma(\tau), \quad \Gamma(1, \tau) = \eta(\tau) \quad \text{for } \tau \in [0, 1]$$

and $\Gamma(\cdot, \tau) : [0, 1] \to V$ is a chain-segment for all $\tau \in [0, 1]$. Note that every chain-segment in this construction is finite in its euclidean length. By Lemma 27, the mapping ϕ is analytically continued along the whole path $\gamma[0, 1]$.

Then the image of E under the mapping φ satisfies

$$\varphi(E) \subset bB^{n+1} \backslash L$$

which yields

$$\varphi(E) = bB^{n+1} \backslash L.$$

Since the mapping ϕ^{-1} is analytically continued on $bB^{n+1} \backslash L$, the mapping $\varphi : E \to bB^{n+1}$ is a CR equivalence. This completes the proof. $\square$

Corollary 29. *Let* D *be a bounded domain with spherical real analytic boundary* bD. *Then there is a natural embedding*

$$Aut(D) \subset Aut\left(B^{n+1}\right).$$

Proof. Let E be the path space of bD pointed at the point $p \in bD$ mod homotopy so that E be a universal covering of bD with a natural CR structure and a natural CR covering map $\psi : E \to bD$. Then we take a biholomorphic mapping ϕ on an open neighborhood U of a point $p \in bD$ such that

$$\phi(U \cap bD) \subset bB^{n+1}.$$

Let L be the Doggaebi variety associated to the mapping ϕ. Then, by Theorem 28, the analytic continuation of the mapping ϕ yields a natural CR equivalence

$$Aut(E) \simeq Aut\left(bB^{n+1} \backslash L\right).$$

Note that

$$Aut\,(D) = Aut(bD) \subset Aut\,(E) \simeq Aut\left(bB^{n+1} \backslash L\right) \subset Aut\left(bB^{n+1}\right) = Aut\left(B^{n+1}\right).$$

This completes the proof. □

Lemma 30. *Let D, D' be bounded domains in $\mathbb{C}^{n+1}$ with spherical real analytic boundaries bD, bD', and $\phi : D \to D'$ be a proper holomorphic mapping. Suppose that there is an open neighborhood U of a point $p \in bD$ such that the mapping ϕ is analytically continued on U. Then the mapping ϕ is analytically continued along any path on bD as a locally biholomorphic mapping so that $\phi : D \to D'$ is locally biholomorphic.*

Proof. We may obtain the result by the boundary regularity of Lemma 16 and the transformation formula for a complex Monge-Ampere equation. Here we may give an independent proof. Note that

$$\phi\,(U \cap bD) \subset bD'$$

so that, by shrinking U, if necessary, the mapping ϕ is biholomorphic on U. By shrinking U, if necessary, we take biholomorphic mappings φ, φ' respectively on $U, \phi\,(U)$ such that

$$\begin{array}{rcl} \varphi\,(U \cap bD) & \subset & bB^{n+1} \\ \varphi'\,(\phi\,(U) \cap bD') & \subset & bB^{n+1}. \end{array}$$

Then there are Doggaebi varieties L, L' on bB^{n+1} such that the inverses $\varphi^{-1}, \varphi'^{-1}$ are analytically continued respectively on $bB^{n+1} \backslash L$ and $bB^{n+1} \backslash L'$ as a local biholomorphic mapping. Note that the composition $\psi = \varphi' \circ \phi \circ \varphi^{-1}$ satisfies the relation

$$\psi\left(\varphi\,(U) \cap bB^{n+1}\right) \subset bB^{n+1}.$$

By abuse of notation, the mapping ψ is an automorphism of bB^{n+1} which comments the following diagram:

$$\begin{tikzcd} bB^{n+1} \backslash L \arrow[r, "\psi"] \arrow[d, "\kappa"] & bB^{n+1} \backslash L' \arrow[d, "\kappa'"] \\ bD \arrow[r, "\phi"] & bD' \end{tikzcd}.$$

where $\kappa : bB^{n+1} \backslash L \to bD$ is the analytic continuation of the mapping φ^{-1} and $\kappa' : bB^{n+1} \backslash L' \to bD'$ is the analytic continuation of the mapping φ'^{-1} on the boundary bB^{n+1}.

We claim

$$L' \subset \psi\,(L)$$

so that the mapping ϕ is analytically continued along any path on bD as a locally biholomorphic mapping. Otherwise, there is a point $q' \in L'$ such that

$$q' \in L' \backslash \psi\,(L)$$

From the proof of First Scaling Lemma, there is a sequence $p'_j \in bD'$ with $p'_j \to p' \in bD'$ and a sequence φ'_j of the analytic continuation of the mapping φ' at the point p'_j with $q'_j \equiv \varphi'_j\left(p'_j\right) \to q' \in L' \backslash \psi\,(L)$ such that there is an open neighborhood V of the point p' such that the mapping φ'_j is biholomorphic on $V \cap D'$ and the sequence

φ_j' converges to the point r uniformly on every compact subset of $V \cap D'$. Since ψ is an automorphism of the unit ball B^{n+1}, we obtain

$$\lambda_j \equiv \kappa \circ \psi^{-1} \circ \varphi_j' : V \cap D' \to D$$

and

$$\phi \circ \lambda_j = id \quad \text{on } V \cap D'.$$

We set

$$p_j \equiv \lambda_j \left(p_j'\right) \in bD.$$

Since bD is compact, there is a subsequence $p_{m_j} \in bD$ and a point $p \in bD$ such that

$$p_{m_j} \to p.$$

Since ϕ is a proper mapping, the mapping ϕ is a globally finite covering so that there is an open neighborhood W of the point p such that

$$\lambda_{m_j} = \phi^{-1} \quad \text{on } \phi\left(W\right) \cap D.$$

Hence we obtain

$$\psi = \varphi_{m_j}' \circ \phi \circ \kappa$$

Since the mapping ψ is an automorphism of the unit ball B^{n+1}, there is a sequence $q_j \in bB^{n+1}$ such that

$$\psi\left(q_j\right) = q_j'.$$

Hence we obtain

$$\kappa\left(q_{m_j}\right) = p_{m_j}.$$

Note that there is a real number $\delta > 0$ such that

$$B\left(r; 2\delta\right) \cap bB^{n+1} \cap \psi\left(L\right) = \emptyset.$$

Passing to a subsequence, if necessary, we may assume

$$q_{m_j} \in \psi^{-1}\left(B\left(r;\delta\right)\right) \cap bB^{n+1}$$

so that there is a point $q \in bB^{n+1} \backslash L$ satisfying

$$q_{m_j} \to q.$$

Hence we set

$$\chi_{m_j} \equiv \psi^{-1} \circ \varphi_{m_j}' \circ \lambda_{m_j}^{-1}$$

and

$$\kappa \circ \chi_{m_j} = id.$$

Finally, we obtain

$$\varphi_{m_j}' = \psi \circ \chi_{m_j} \circ \lambda_{m_j}.$$

Note that there is a real number $\delta > 0$ that the mappings λ_{m_j} and the inverse mappings $\lambda_{m_j}^{-1}$ are analytically continued respectively on

$$B\left(p_{m_j}'; \delta\right) \quad \text{and} \quad B\left(p_{m_j}; \delta\right),$$

and the mappings χ_{m_j} and the inverse mappings $\chi_{m_j}^{-1}$ are analytically continued respectively on

$$B\left(p_{m_j};\delta\right) \quad \text{and} \quad B\left(q_{m_j};\delta\right),$$

and the mapping ψ and the inverse mapping ψ^{-1} are analytically continued respectively on

$$B\left(q_{m_j};\delta\right) \quad \text{and} \quad B\left(q'_{m_j};\delta\right),$$

Hence the mapping φ'_{m_j} and the inverse mappings $\varphi'^{-1}_{m_j}=\kappa'$ are analytically continued respectively on

$$B\left(p'_{m_j};\delta\right) \quad \text{and} \quad B\left(q'_{m_j};\delta\right),$$

as a locally biholomorphic mapping. Since $q'_{m_j}\rightarrow q'$, the mapping κ' is analytically continued to the point q' as a locally biholomorphic mapping so that

$$q'\in bB^{n+1}\backslash L'.$$

This is a contradiction so that

$$L'\subset\psi\left(L\right).$$

This completes the proof. □

2. Bounded Domains with Spherical Boundaries

2.1. **Differentiable spherical boundary.** Let D be a domain in $\mathbb{C}^{n+1}, n\geq 1$, with a differentiable boundary bD. The boundary bD shall be called spherical if, for each point $p\in bD$, there is a connected open neighborhood U of the point p and a biholomorphic mapping ϕ on $U\cap D$ such that

$$\phi\in H\left(U\cap D\right)\cap C^1\left(U\cap\overline{D}\right), \quad \phi\left(U\cap bD\right)\subset bB^{n+1} \tag{13}$$

and the induced mapping $\phi: U\cap bD\rightarrow bB^{n+1}$ is CR diffeomorphic.

Lemma 31. *Let U be a connected open neighborhood of a point $p\in bB^{n+1}$ and ϕ be a biholomorphic mapping on $U\cap B^{n+1}$ such that*

$$\phi\in H\left(U\cap B^{n+1}\right)\cap C^1\left(U\cap\overline{B^{n+1}}\right), \quad \phi\left(U\cap bB^{n+1}\right)\subset bB^{n+1}$$

and the induced mapping $\phi: U\cap bB^{n+1}\rightarrow bB^{n+1}$ is CR diffeomorphic. Then the mapping ϕ is analytically continued to an automorphism of the unit ball B^{n+1}.

Proof. By Lemma 13, the mapping ϕ is biholomorphic on U, if necessary, shrinking U. Then, by Lemma 17, we obtain the desired result. This completes the proof. □

The chain on bB^{n+1} is defined to be the intersection on bB^{n+1} by a complex line. The family of chains on bB^{n+1} leaves invariant under the action of biholomorphic automorphisms of B^{n+1}. We define the chain on the spherical differentiable boundary bD of a domain D to be the inverse image of the chain on bB^{n+1} under the mapping 13. By Lemma 31, the chain on the spherical differentiable boundary bD is well defined so that a chain of a spherical differentiable boundary is mapped to a chain of another spherical differentiable boundary under any induced CR diffeomorphism.

Lemma 32. *Let D be a domain in $\mathbb{C}^{n+1}$ with spherical differentiable boundary bD. Suppose that there is a connected open neighborhood U of a point $p \in bD$ and a biholomorphic mapping ϕ on $U \cap D$ such that*

$$\phi \in H(U \cap D) \cap C^1\left(U \cap \overline{D}\right), \quad \phi(U \cap bD) \subset bB^{n+1}.$$

Then the mapping ϕ is analytically continued along any path on bD as a local biholomorphic mapping.

Proof. Suppose that the assertion is not true. Then there would be a path $\gamma : [0,1] \to bD$ such that $\gamma(0) \in U \cap bD$ and the germ of a biholomorphic mapping ϕ at the point $\gamma(0)$ is analytically continued along the subpath $\gamma[0,\tau]$ with all $\tau < 1$ as a local biholomorphic mapping, but not the whole path $\gamma[0,1]$.

Thus we reduce the proof to a local problem near the point $\gamma(1) \in bD$. Then, by the definition, there is a connected open neighborhood V of the point $\gamma(1)$ and a biholomorphic mapping φ on $V \cap D$ such that

$$\varphi \in H(V \cap D) \cap C^1\left(V \cap \overline{D}\right), \quad \varphi(V \cap bD) \subset bB^{n+1}$$

and the induced mapping $\varphi : U \cap bD \to bB^{n+1}$ is CR diffeomorphic.

Then we consider the mapping

$$\phi \circ \varphi^{-1} \in H(\varphi(V \cap D)) \cap C^1\left(\varphi\left(V \cap \overline{D}\right)\right), \quad \phi \circ \varphi^{-1}(\varphi(V \cap bD)) \subset bB^{n+1}$$

and the curve

$$\varphi \circ \gamma[0,1] \cap \varphi(V \cap bD) \subset bB^{n+1}.$$

By Lemma 13, the remaining part of the proof repeats the proof of Lemma 18. This completes the proof. □

Lemma 33 (First Dogginal Lemma). *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical differentiable boundary bD and ϕ be a biholomorphic mapping on $U \cap D$ for a connected open neighborhood U of a point $p \in bB^{n+1}$ satisfying*

$$\phi \in H\left(U \cap B^{n+1}\right) \cap C^1\left(U \cap \overline{B^{n+1}}\right), \quad \phi\left(U \cap bB^{n+1}\right) \subset bD.$$

Suppose that there is an injective path $\gamma : [0,1] \to bB^{n+1}$ such that $\gamma[0,1] \subset bB^{n+1}$ is a chain-segment satisfying

$$\gamma(0) \in U \cap bB^{n+1}$$

and the mapping ϕ is analytically continued along the subpath $\gamma[0,\tau]$ for all $\tau < 1$, but not the whole path $\gamma[0,1]$ as a local biholomorphic mapping. Let π be the complex line containing the chain-segment $\gamma[0,1]$. Then there is an open neighborhood V along the path $\gamma[0,1]$ such that

1. $\gamma[0,\tau] \subset V$ *for all* $\tau < 1$,
2. $bV \cap \pi \cap B(\gamma(1);\delta)$ *is an angle for a sufficient small* $\delta > 0$, *which contains the chain-segment* $\gamma[0,1]$,
3. $bV \cap bB^{n+1} \cap B(\gamma(1);\delta)$ *is paraboloid for a sufficiently small* $\delta > 0$, *which smoothly touches the complex tangent hyperplane at the point* $\gamma(1)$,
4. *the mapping ϕ is analytically continued on $V \cap B^{n+1}$ as a local biholomorphic mapping*

$$\phi \in H\left(V \cap B^{n+1}\right) \cap C^1\left(V \cap \overline{B^{n+1}}\right).$$

Proof. By the analytic continuation of the mapping ϕ along the subpath $\gamma[0,\tau]$ for all $\tau < 1$, there is a path $\phi \circ \gamma : [0,1) \to bD$. Then we consider the following sequences

$$\begin{aligned} p_j &= \gamma\left(1-\frac{1}{j}\right), \quad \text{for } j \in \mathbb{N}^+, \\ p_j' &= \phi \circ \gamma\left(1-\frac{1}{j}\right), \quad \text{for } j \in \mathbb{N}^+. \end{aligned}$$

Since bD is compact, there is a subsequence p'_{m_j} and a point $p' \in bD$ such that

$$p'_{m_j} \to p'.$$

Thus we reduce the proof to a local problem near the point $p' \in bD$. Then, by the definition, there is a connected open neighborhood W of the point p' and a biholomorphic mapping φ on $W \cap D$ such that

$$\varphi \in H\left(W \cap D\right) \cap C^1\left(W \cap \overline{D}\right), \quad \varphi\left(W \cap bD\right) \subset bB^{n+1}$$

and the induced mapping $\varphi : U \cap bD \to bB^{n+1}$ is CR diffeomorphic.

Then we consider the mapping

$$\phi^{-1} \circ \varphi^{-1} \in H\left(\varphi\left(W \cap D\right)\right) \cap C^1\left(\varphi\left(W \cap \overline{D}\right)\right), \quad \phi^{-1} \circ \varphi^{-1}\left(\varphi\left(W \cap bD\right)\right) \subset bB^{n+1}.$$

By Lemma 13, the remaining part of the proof repeats the proof of Lemma 24. This completes the proof. □

By Lemma 31 and Lemma 33, we obtain the following result by the same argument of the proof of Theorem 28.

Theorem 34. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical differentiable boundary bD. Suppose that there is a biholomorphic mapping ϕ on $U \cap D$ for a connected open neighborhood U of a point $p \in bD$ satisfying*

$$\phi\left(U \cap bD\right) \subset bB^{n+1}.$$

Let E be the path space of bD pointed at the point $p \in bD$ mod homotopy so that E be a universal covering of bD with a natural CR structure and a natural CR covering map $\psi : E \to bD$. Then there is a unique CR equivalence $\varphi : E \to bB^{n+1}\backslash L$ commuting the diagram

$$\begin{tikzcd} E \arrow[d, "\psi"'] \arrow[dr, "\varphi"] & \\ bD \arrow[r, "\phi"] & bB^{n+1}\backslash L \end{tikzcd}$$

where $L \subset bB^{n+1}$ is the Doggaebi variety associated to the mapping ϕ.

2.2. **Second Dogginal Lemma.**

We have examined the analytic continuation of a biholomorphic mapping ϕ from the spherical differentiable boundary bD of a bounded domain D into the boundary bB^{n+1} of the unit ball B^{n+1}. From now on, we shall examine the analytic continuation of the mapping ϕ into the domain D.

Lemma 35. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical differentiable boundary bD. Suppose that there is a biholomorphic mapping ϕ on a connected open neighborhood U of a point $p \in bD$ satisfying*

$$\phi\left(U \cap bD\right) \subset bB^{n+1}.$$

Then the inverse mapping ϕ^{-1} is analytically continued to a locally biholomorphic mapping from B^{n+1} into D.

Proof. By Theorem 34, there is a Doggaebi variety L on bB^{n+1} such that the mapping ϕ^{-1} is uniquely analytically continued on the set $bB^{n+1}\backslash L$. Then the mapping ϕ^{-1} is analytically continued to a holomorphic mapping on the unit ball B^{n+1}. Hence we have a holomorphic mapping $\varphi : B^{n+1} \to D$ which extends to the mapping $\phi^{-1} : bB^{n+1}\backslash L \to bD$. Note that the zero set of the determinant of the Jacobian matrix φ' of the mapping φ cannot be located on the set L on the boundary bB^{n+1}. Thus the mapping $\varphi : B^{n+1} \to D$ is locally biholomorphic. This completes the proof. □

Lemma 36. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical differentiable boundary bD and ϕ be a biholomorphic mapping on an open neighborhood U of a point $r \in bD$ satisfying*

$$\phi \in H\left(U \cap D\right) \cap C^1\left(U \cap \overline{D}\right) \quad \text{and} \quad \phi\left(U \cap bD\right) \subset bB^{n+1}.$$

Let L be the Doggaebi variety associated to the mapping ϕ and $\varphi : B^{n+1} \to D$ be a locally biholomorphic mapping to be an analytic continuation of the mapping ϕ^{-1}. Suppose that there is a line segment $\gamma : [0,1] \to D$ with

$$p = \gamma\left(1\right) \quad \text{and} \quad p_j = \gamma\left(1 - \frac{1}{j}\right)$$

and the germ of a locally biholomorphic mapping $\phi = \varphi^{-1}$ at the point $p_1 \equiv \gamma\left(0\right)$ is analytically continued along the subpath $\gamma[0,\tau]$ with all $\tau < 1$ as a locally biholomorphic mapping. Let $p'_j \equiv \phi\left(p_j\right) \in B^{n+1}$ be the sequence obtained by the analytic continuation of the mapping $\phi = \varphi^{-1}$ along the path $\gamma[0,1)$. Then there is a subsequence p'_{m_j} and a point $p' \in L \subset bB^{n+1}$ such that the subsequence p'_{m_j} converges to the point p' to a direction transversal to the complex tangent hyperplane at the point $p' \in bB^{n+1}$.

Proof. Since the closed ball $\overline{B^{n+1}}$ is compact, there is a subsequence p'_{m_j} which converges to a point $p' \in \overline{B^{n+1}}$. Since the mapping φ is locally biholomorphic on B^{n+1}, the point p' should be on the boundary bB^{n+1}. Further, since $\varphi = \phi^{-1} : bB^{n+1}\backslash L \to bD$ is locally biholomorphic, we obtain

$$p'_{m_j} \to p' \in L.$$

Let π_p be the complex line containing the line segment $\gamma[0,1]$. Since $p'_{m_j} \equiv \phi\left(p_{m_j}\right)$, the analytic continuation of the mapping $\phi = \varphi^{-1}$ on the complex line π_p yields a complex curve $\varphi^{-1}\left(\pi_p \cap D\right)$ which touches the point $p' \in L \subset bB^{n+1}$. Since the mapping $\varphi : B^{n+1} \to D$ is locally biholomorphic up to the boundary subset $bB^{n+1}\backslash L$ and the boundary bB^{n+1} is strongly pseudoconvex, the extension

$$\varphi^{-1}\left(\pi_p \cap \overline{D}\right)$$

of the complex curve $\varphi^{-1}\left(\pi_p \cap D\right)$ is transversal to the boundary bB^{n+1} near the point $p' \in L \subset bB^{n+1}$.

We claim that the extension $\varphi^{-1}\left(\pi_p \cap \overline{D}\right)$ is transversal to the complex tangent hyperplane at the point $p' \in L$. We take a path $\lambda : [0,1] \to bB^{n+1}$ such that

$p' = \lambda(1)$ and

$$\lambda[0,1] \subset \overline{\varphi^{-1}\left(\pi_p \cap bD\right)} \cap bB^{n+1}.$$

Note that there are finitely many closed paths γ_j on bD such that

$$\pi_p \cap bD = \bigcup_j \gamma_j.$$

Since bD is strongly pseudoconvex, each path γ_j is transversal to the complex tangent hyperplane on bD at each point.

We consider the following sequence

$$q_j = \varphi \circ \lambda \left(1 - \frac{1}{j}\right) \in \pi_p \cap bD, \quad q_j' = \lambda\left(1 - \frac{1}{j}\right) \quad \text{for } j \in \mathbb{N}^+$$

such that

$$q_j' \to p' \quad \text{as } j \to \infty.$$

Since the set $\pi_p \cap bD$ is compact, there is a subsequence q_{m_j} and a point $q \in \pi_p \cap bD$ such that

$$q_{m_j} \to q \quad \text{as } j \to \infty.$$

Let ϕ_{m_j} be the germ of the analytic continuation of the mapping $\phi = \varphi^{-1}$ at the point q_{m_j}. By Theorem 18 and Lemma 14, we may assume that there is an open neighborhood W of the point q such that the mapping ϕ_{m_j} is holomorphic on $W \cap D$.

Since bD is spherical, there is a biholomorphic mapping ψ on W, if necessary, shrinking W, such that

$$\psi \in H\left(W \cap D\right) \cap C^1\left(W \cap \overline{D}\right) \quad \text{and} \quad \psi\left(W \cap bD\right) \subset bB^{n+1}.$$

We may assume that

$$q'' \equiv \psi\left(q\right) \neq p' \quad \text{and} \quad q_{m_j}'' \equiv \psi\left(q_{m_j}\right).$$

Then, by Lemma 17, the composition

$$\eta_j \equiv \phi_{m_j} \circ \psi^{-1} \in Aut\left(B^{n+1}\right)$$

are automorphisms of the unit ball B^{n+1}. Note that we have the condition

$$\eta_j\left(q_{m_j}''\right) = q_{m_j}' \to p'$$

and the sequence η_j converges to the point p' uniformly on every compact subset of B^{n+1}, if necessary, passing to a subsequence. Otherwise, $p' \notin L$.

Let π_j' be the complex line passing through the two points p' and q_{m_j}', and δ_j' be the analytic disk

$$\delta_j' = \pi_j' \cap B^{n+1}.$$

The area $\left|\eta_j^{-1}\left(\delta_j'\right)\right|$ of the analytic disk $\eta_j^{-1}\left(\delta_j'\right)$ is bounded from the below. Otherwise, we would have $p' \notin L$. Then we decompose the mapping η_j as follows

$$\eta_j = \mu_j \circ \nu_j$$

where

$$\mu_j \in Aut_{q_{m_j}'}\left(bB^{n+1}\right), \quad \nu_j \in Aut_{p'}\left(bB^{n+1}\right).$$

Then, by First and Second Scaling Lemmas, we take a sequence of local automorphisms $\sigma_j \in Aut_{q'_{m_j}}\left(bB^{n+1}\right)$ defined by the normalizing parameters

$$U_{\sigma_j} = id_{n\times n}, \quad \rho_{\sigma_j} = \rho_{\mu_j}, \quad a_{\sigma_j} = 0, \quad r_{\sigma_j} = 0.$$

Then the following sequence

$$\beta_j = \sigma_j^{-1} \circ \eta_j$$

converges to an automorphism of the unit ball B^{n+1}, if necessary, passing to a subsequence.

Let π_j be the complex line containing the tangent vector of the path

$$\psi\left(\varphi \circ \lambda[0,1] \cap W\right) \subset bB^{n+1}$$

at the point q''_{m_j}, and δ_j be the analytic disk

$$\delta_j = \pi_j \cap B^{n+1}.$$

Clearly, the area $|\delta_j|$ of the analytic disk δ_j is bounded from the below. Note that, by the mapping $\sigma_j \in Aut_{q'_{m_j}}\left(bB^{n+1}\right)$, the area $\left|\sigma_j^{-1}\left(\delta''_j\right)\right|$ of the analytic disk $\sigma_j\left(\delta''_j\right)$ is bounded from the below whenever the area $\left|\delta''_j\right|$ of the analytic disk δ''_j is bounded from the below. Thus the area $|\eta_j\left(\delta_j\right)|$ of the analytic disk

$$\eta_j\left(\delta_j\right) = \sigma_j \circ \beta_j\left(\delta_j\right)$$

is bounded from the below. Since the analytic disk $\eta_j\left(\delta_j\right)$ is the intersection of B^{n+1} and the complex line containing the tangent vector of the path $\lambda[0,1]$ at the point

$$q'_{m_j} = \lambda\left(1 - \frac{1}{m_j}\right).$$

Thus the path $\lambda[0,1]$ is transversal to the complex tangent hyperplane at the point $p' = \lambda(1)$. Therefore, the point p'_j approaches to the point $p' \in L$ to a direction transversal to the complex tangent hyperplane at the point $p' \in bB^{n+1}$. This completes the proof. □

Lemma 37 (Second Dogginal Lemma)**.** *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical differentiable boundary bD and ϕ be a biholomorphic mapping on an open neighborhood U of a point $r \in bD$ satisfying*

$$\phi \in H\left(U \cap D\right) \cap C^1\left(U \cap \overline{D}\right) \quad \textit{and} \quad \phi\left(U \cap bD\right) \subset bB^{n+1}.$$

Let L be the Doggaebi variety associated to the mapping ϕ and $\varphi : B^{n+1} \rightarrow D$ be a locally biholomorphic mapping to be an analytic continuation of the mapping ϕ^{-1}. Suppose that there is a line segment $\gamma : [0,1] \rightarrow D$ with

$$p = \gamma(1) \quad \textit{and} \quad p_j = \gamma\left(1 - \frac{1}{j}\right)$$

and the germ of a locally biholomorphic mapping $\phi = \varphi^{-1}$ at the point $p_1 \equiv \gamma(0)$ is analytically continued along the subpath $\gamma[0,\tau]$ with all $\tau < 1$ as a locally biholomorphic mapping. Let $p'_j \equiv \phi(p_j) \in B^{n+1}$ be the sequence obtained by the analytic continuation of the mapping $\phi = \varphi^{-1}$ along the path $\gamma[0,1)$. Suppose that there is a point $p' \in L \subset bB^{n+1}$ such that the sequence p'_j converges to the point p' to a direction transversal to the complex tangent hyperplane at the point $p' \in bB^{n+1}$.

Let π_p be the complex line containing the line segment $\gamma[0,1]$. Then there is a distinguished complex hyperplane $H_p \subset T_p\mathbb{C}^{n+1}$ at the point $p \in D$ satisfying

$$T_p\mathbb{C}^{n+1} = H_p \oplus \pi_p$$

and an open neighborhood V of the line segment $\gamma[0,1)$ such that

1. *$bV \cap \pi_p \cap B(p;\delta)$ is an angle for a sufficient small $\delta > 0$, which contains the path $\gamma[0,1]$,*
2. *$bV \cap R_p \cap B(p;\delta)$ is a slanted paraboloid for a sufficiently small $\delta > 0$, which smoothly touches the complex hyperplane H_p at the point p,*
3. *the germ of a locally biholomorphic mapping $\phi = \varphi^{-1}$ at the point $p_1 \equiv \gamma(0)$ is analytically continued on V as a locally biholomorphic mapping,*

where R_p is a real hyperplane containing the line segment $\gamma[0,1]$ and

$$H_p \subset R_p.$$

Proof. We take a sequence of automorphisms $\phi_j \in Aut\left(B^{n+1}\right)$ such that

$$\phi_j\left(p_1'\right) = p_j' \to p' \in L.$$

Then the composition $\tau_j \equiv \varphi \circ \phi_j : B^{n+1} \to D$ forms a normal family so that there is a subsequence $\tau_{m_j} \equiv \varphi \circ \phi_{m_j}$ which converges to a holomorphic mapping $\tau : B^{n+1} \to D$ uniformly on every compact subset of B^{n+1}.

We claim that τ is a locally biholomorphic mapping. We take a point $q \in bB^{n+1}\backslash L$ and $q'_{m_j} = \tau_{m_j}(q) \in bD$ such that

$$q'_{m_j} = \tau_{m_j}(q) \to q' \in bD,$$

if necessary, passing to a subsequence. Since bD is spherical, there is an open neighborhood W of the point q' and a biholomorphic mapping ψ on $W \cap D$ satisfying

$$\psi \in H(W \cap D) \cap C^1\left(W \cap \overline{D}\right) \quad \text{and} \quad \psi(W \cap bD) \subset bB^{n+1}.$$

Then we consider the composition $\chi_{m_j} \equiv \psi \circ \varphi \circ \phi_{m_j}$ so that, by Lemma 17 and by abuse of notation,

$$\chi_{m_j} \in Aut\left(B^{n+1}\right).$$

Suppose that the assertion is not true. Then the sequence χ_{m_j} converges to a boundary point $\psi(q') \in bB^{n+1}$ uniformly on every compact subset of B^{n+1}, if necessary, passing to a subsequence. Hence the sequence $\tau_{m_j} \equiv \varphi \circ \phi_{m_j}$ would converge to a boundary point $q' \in bD$ uniformly on every compact subset of B^{n+1}. By the way, note that

$$\tau_{m_j}\left(p_1'\right) = p_{m_j} \to p \in D.$$

This is a contradiction. Hence the holomorphic mapping τ is locally biholomorphic. Thus there is a real number $\delta > 0$ such that the mapping τ_{m_j} and the inverse $\tau_{m_j}^{-1}$ are analytically continued to a biholomorphic mapping respectively on

$$B\left(p_{m_j};\delta\right), \quad B\left(p_1';\delta\right).$$

Let ε_j be the euclidean length between the two points p' and p_j'. By First Scaling Lemma, there is a sequence of automorphisms

$$\sigma_j \in Aut_{p'}\left(bB^{n+1}\right)$$

such that, for a subsequence ϕ_{m_j},

$$U_{\sigma_j} = id_{n\times n}, \quad \rho_{\sigma_j} = \varepsilon_{m_j}, \quad a_{\sigma_j} = 0, \quad r_{\sigma_j} = 0$$

and the composition

$$\eta_j \equiv \sigma_j^{-1} \circ \phi_{m_j}$$

uniformly converges to an automorphism of the unit ball B^{n+1}. Then there is a real number $\delta > 0$ such that the mapping η_j and the inverse η_j^{-1} are analytically continued respectively on

$$B\left(p_1'; \delta\right), \quad B\left(p_{m_j}''; \delta\right)$$

where

$$p_{m_j}'' = \eta_j\left(p_{m_j}'\right).$$

Then we obtain

$$\varphi = \tau_{m_j} \circ \eta_j^{-1} \circ \sigma_j^{-1}.$$

Thus the mapping $\phi = \varphi^{-1}$ is analytically continued onto the open set

$$\tau_{m_j} \circ \eta_j^{-1}\left(\sigma_j\left(B\left(p_{m_j}''; \delta\right)\right)\right)$$

which is centered at the point p_{m_j}. For a canonical normalizing mapping $\mu_{p'}$, we obtain

$$\sigma_j' \equiv \mu_{p'} \circ \sigma_j \circ \mu_{p'}^{-1} : \left\{ \begin{array}{l} z^* = \sqrt{\varepsilon_{m_j}} z \\ w^* = \varepsilon_{m_j} w \end{array} \right. .$$

We set

$$\mu_{p'}\left(p_{m_j}''\right) \to p''' \in \mu_{p'}\left(B^{n+1}\right)$$

so that, shrinking $\delta > 0$, if necessary, the mapping $\phi = \varphi^{-1}$ is analytically continued onto the open set

$$\tau_{m_j} \circ \eta_j^{-1} \circ \mu_{p'}\left(\sigma_j'\left(B\left(p'''; \delta\right)\right)\right)$$

which is centered at the point p_{m_j}.

Since ε_{m_j} is the euclidean distance between the two points p and p_{m_j}, the analytically continued region of the mapping $\phi = \varphi^{-1}$ along the line segment $\gamma[0,1] \subset D$ contains an open set along the line segment $\gamma[0,1]$ which touches to the point $p = \gamma(1)$ by an edge shape on the complex line π_p and by a slanted paraboloid shape on the real hyperplane R_p. This completes the proof. □

Lemma 38. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical differentiable boundary bD and ϕ be a biholomorphic mapping on an open neighborhood U of a point $r \in bD$ satisfying*

$$\phi \in H\left(U \cap D\right) \cap C^1\left(U \cap \overline{D}\right) \quad \text{and} \quad \phi\left(U \cap bD\right) \subset bB^{n+1}.$$

Let L be the Doggaebi variety associated to the mapping ϕ and $\varphi : B^{n+1} \to D$ be a locally biholomorphic mapping to be an analytic continuation of the mapping ϕ^{-1}. Let H_p be the complex hyperplane at the point $p \in D$ in Lemma 37 and π_p be a complex line satisfying

$$T_p\mathbb{C}^{n+1} = H_p \oplus \pi_p.$$

Suppose that, along a line segment $\gamma : [0,1] \to \pi_p \cap D$ *with* $p = \gamma(1)$, *the germ of a locally biholomorphic mapping* $\phi = \varphi^{-1}$ *at the point* $\gamma(0)$ *is analytically continued along the subpath* $\gamma[0,\tau]$ *with all* $\tau < 1$ *as a locally biholomorphic mapping such that the limit point*

$$\lim_{\tau \to 1} \phi \circ \gamma(\tau)$$

is the point $p' \in L \subset bB^{n+1}$ *in Lemma 37. Then there is an open neighborhood* V *of the line segment* $\gamma[0,1)$ *such that*

1. $bV \cap \pi_p \cap B(p;\delta)$ *is an angle for a sufficient small* $\delta > 0$, *which contains the path* $\gamma[0,1]$,
2. $bV \cap R_p \cap B(p;\delta)$ *is a slanted paraboloid for a sufficiently small* $\delta > 0$, *which smoothly touches the complex hyperplane* H_p *at the point* p,
3. *the germ of a locally biholomorphic mapping* $\phi = \varphi^{-1}$ *at the point* $\gamma(0)$ *is analytically continued on* V *as a locally biholomorphic mapping,*

where R_p *is a real hyperplane containing the line segment* $\gamma[0,1]$ *and*

$$H_p \subset R_p.$$

Proof. By Lemma 36, $\phi \circ \gamma(\tau)$ approaches to the point $p' \in L$ to a direction transversal to the complex tangent hyperplane at the point p' as $\tau \to 1$. Then there is a complex hyperplane H_p' satisfying the conditions in Lemma 37.

Note that the complex hyperplane H_p is determined by the mapping $\varphi : B^{n+1} \to D$ and the complex tangent hyperplane $H_{p'}$ at the point $p' \in L \subset bB^{n+1}$, but H_p does not depend on the approaching direction of $\phi \circ \gamma(\tau) \to p'$ as $\tau \to 1$. Thus $H_p' = H_p$. This completes the proof. □

2.3. **Biholomorphic equivalence.**

Lemma 39. *Let* D *be a bounded domain in* $\mathbb{C}^{n+1}$ *with spherical differentiable boundary* bD *and* G *be the path space of* D *mod homotopic relation so that* G *is a universal covering Riemann domain with a natural complex structure and a natural holomorphic covering map* $\kappa : G \to D$. *Suppose that there is a biholomorphic mapping*

$$\phi \in H(U \cap D) \cap C^1\left(U \cap \overline{D}\right)$$

for a connected open neighborhood U *of a point* $p \in bD$ *satisfying*

$$\phi(U \cap bD) \subset bB^{n+1}.$$

Then the analytic continuation of the inverse mapping ϕ^{-1} *is lifted to a holomorphic mapping* ψ *from* B^{n+1} *onto* G *such that* $\psi : B^{n+1} \to G$ *is a proper locally biholomorphic mapping, i.e., a finite covering of* G, *satisfying the following relation:*

$$\begin{tikzcd} & G \arrow[d, "\kappa"] \\ B^{n+1} \arrow[ur, "\psi"] \arrow[r, "\varphi"] & D \end{tikzcd} .$$

Proof. By Theorem 35, there is a locally biholomorphic mapping

$$\varphi : B^{n+1} \to D$$

such that φ is the analytic continuation of the mapping ϕ^{-1} and

$$\varphi = \phi^{-1} \quad \text{on} \quad bB^{n+1} \backslash L$$

where L is the Doggaebi variety associated to the mapping ϕ.

A piecewise line segment curve is a path $\gamma : [0,1] \to D$ consisting of finitely many line segments. Fix a point $b \in U \cap D$. We shall show that the mapping ϕ is analytically continued along any piecewise line segment curve. Suppose that $\gamma : [0,1] \to D$ be a piecewise line segment curve with $\gamma(0) = b$ such that the germ of the mapping ϕ at the point $\gamma(0)$ is analytically continued along all subpath $\gamma[0,\tau]$ with $\tau < 1$, but not the whole path $\gamma[0,1]$. Let π_p be a complex line satisfying, for sufficiently small $\varepsilon > 0$,

$$\gamma[1-\varepsilon, 1] \subset \pi_p.$$

Then, by Second Dogginal Lemma, there is a distinguished complex hyperplane H_p at the point $p = \gamma(1)$ satisfying

$$T_p\mathbb{C}^{n+1} = H_p \oplus \pi_p$$

and an open neighborhood V containing the line segment $\gamma[0,\tau]$ with all $\tau < 1$ such that

1. $bV \cap \pi_p \cap B(p;\varepsilon)$ is an angle for a sufficient small $\varepsilon > 0$, which contains the path $\gamma[0,1]$,
2. $bV \cap R_p \cap B(p;\varepsilon)$ is a slanted paraboloid for a sufficiently small $\varepsilon > 0$, which smoothly touches the complex hyperplane H_p at the point p,
3. the germ of a locally biholomorphic mapping $\phi = \varphi^{-1}$ at the point $\gamma(0)$ is analytically continued on V as a locally biholomorphic mapping,

where R_p is a real hyperplane satisfying, for sufficiently small $\varepsilon > 0$,

$$\gamma[1-\varepsilon, 1] \subset R_p \quad \text{and} \quad H_p \subset R_p.$$

Thus we can find a sequence of points $p_j \in V \cap B(p;\varepsilon)$ and a sequence of germs ϕ_j of the analytic continuation of the mapping ϕ at the point p_j such that the sequence p_j converges to the point p to a direction tangential to the complex hyperplane H_p at the point p. We set

$$p'_j = \phi_j(p_j) \in B^{n+1}.$$

Then there is a point $p' \in L \subset bB^{n+1}$ such that $p'_j \to p'$ to a direction tangential to the complex tangent hyperplane at the point $p' \in bB^{n+1}$. We take a sequence of automorphisms $\varphi_j \in Aut\left(B^{n+1}\right)$ such that

$$\varphi_j(p'_1) = p'_j \to p' \in L.$$

Then the composition $\tau_j \equiv \varphi \circ \varphi_j : B^{n+1} \to D$ forms a normal family so that there is a subsequence $\tau_{m_j} \equiv \varphi \circ \varphi_{m_j}$ which converges to a holomorphic mapping $\tau : B^{n+1} \to D$ uniformly on every compact subset of B^{n+1}.

We claim that τ is a locally biholomorphic mapping. We take a point $q \in bB^{n+1} \backslash L$ and $q'_{m_j} = \tau_{m_j}(q) \in bD$ such that

$$q'_{m_j} = \tau_{m_j}(q) \to q' \in bD,$$

if necessary, passing to a subsequence. Since bD is spherical, there is an open neighborhood W of the point q' and a biholomorphic mapping ψ on $W \cap D$ satisfying

$$\psi \in H(W \cap D) \cap C^1\left(W \cap \overline{D}\right) \quad \text{and} \quad \psi(W \cap bD) \subset bB^{n+1}.$$

Then we consider the composition $\chi_{m_j} \equiv \psi \circ \varphi \circ \varphi_{m_j}$ so that, by Lemma 17 and by abuse of notation,

$$\chi_{m_j} \in Aut\left(B^{n+1}\right).$$

Suppose that the assertion is not true. Then the sequence χ_{m_j} converges to a boundary point $\psi\left(q'\right) \in bB^{n+1}$ uniformly on every compact subset of B^{n+1}, if necessary, passing to a subsequence. Hence the sequence $\tau_{m_j} \equiv \varphi \circ \varphi_{m_j}$ would converge to a boundary point $q' \in bD$ uniformly on every compact subset of B^{n+1}. By the way, note that

$$\tau_{m_j}\left(p'_1\right) = p_{m_j} \to p \in D.$$

This is a contradiction. Hence the holomorphic mapping τ is locally biholomorphic. Thus there is a real number $\delta > 0$ such that the mapping τ_{m_j} and the inverse $\tau_{m_j}^{-1}$ are analytically continued to a biholomorphic mapping respectively on

$$B\left(p_{m_j};\delta\right), \quad B\left(p'_1;\delta\right).$$

Let ε_j be the euclidean length between the two points p' and p'_j. By Second Scaling Lemma, there is a sequence of automorphisms

$$\sigma_j \in Aut_{p'}\left(bB^{n+1}\right)$$

such that, for a subsequence φ_{m_j},

$$U_{\sigma_j} = id_{n\times n}, \quad \rho_{\sigma_j} = \varepsilon_{m_j}^2, \quad a_{\sigma_j} = 0, \quad r_{\sigma_j} = 0$$

and the composition

$$\eta_j \equiv \sigma_j^{-1} \circ \varphi_{m_j}$$

uniformly converges to an automorphism of the unit ball B^{n+1}. Then there is a real number $\delta > 0$ such that the mapping η_j and the inverse η_j^{-1} are analytically continued respectively on

$$B\left(p'_1;\delta\right), \quad B\left(p''_{m_j};\delta\right)$$

where

$$p''_{m_j} = \eta_j\left(p'_{m_j}\right).$$

Then we obtain

$$\varphi = \tau_{m_j} \circ \eta_j^{-1} \circ \sigma_j^{-1}.$$

Thus the mapping $\phi = \varphi^{-1}$ is analytically continued onto the open set

$$\tau_{m_j} \circ \eta_j^{-1}\left(\sigma_j\left(B\left(p''_{m_j};\delta\right)\right)\right)$$

which is centered at the point p_{m_j}. For a canonical normalizing mapping $\mu_{p'}$, we obtain

$$\sigma'_j \equiv \mu_{p'} \circ \sigma_j \circ \mu_{p'}^{-1} : \begin{cases} z^* = \varepsilon_{m_j} z \\ w^* = \varepsilon_{m_j}^2 w \end{cases}.$$

We set

$$\mu_{p'}\left(p''_{m_j}\right) \to p''' \in \mu_{p'}\left(B^{n+1}\right)$$

so that, shrinking $\delta > 0$, if necessary, the mapping $\phi = \varphi^{-1}$ is analytically continued onto the open set

$$\tau_{m_j} \circ \eta_j^{-1} \circ \mu_{p'} \left(\sigma_j' \left(B \left(p'''; \delta \right) \right) \right)$$

which is centered at the point p_{m_j}.

Since the sequence p_{m_j} converges to p to a direction tangential to the complex hyperplane H_p at the point p, and ε_{m_j} is the euclidean distance between the two points p and p_{m_j}, the analytically continued region of the mapping $\phi = \varphi^{-1}$ along the sequence $p_{m_j} \in V \cap D$ contains an open set along the converging direction of the sequence p_{m_j} by an edge shape tangential to the complex hyperplane H_p at the point p and a $\sqrt{|x|}$ curve shape normal to H_p.

Thus there is a complex line π_p' satisfying

$$\pi_p' \subset H_p \quad \text{and} \quad p \in \pi_p',$$

and an open neighborhood W of the point p such that the mapping ϕ is analytically continued on

$$W \cap \pi_p'.$$

Then we take a line segment $\gamma : [0,1] \to \pi_p' \subset H_p$ such that

$$\gamma : [0,1) \subset W \cap \left(\pi_p' \backslash p \right) \quad \text{and} \quad \gamma(1) = p.$$

Note that

$$\lim_{\tau \to 1} \phi \circ \gamma(\tau) = p' \in L \subset bB^{n+1}.$$

By Second Dogginal Lemma, the distinguished complex tangent hyperplane H_p satisfies

$$T_p \mathbb{C}^{n+1} = H_p \oplus \pi_p'$$

so that

$$\pi_p' \cap H_p = \{p\}.$$

This is a contradiction so that the germ of the mapping ϕ at the point $b \in U \cap D$ is analytically continued along the whole path $\gamma[0,1]$. Hence the mapping ϕ is analytically continued along any piecewise line segment curve on D as a locally biholomorphic mapping.

We claim that the mapping $\phi : U \cap D \to B^{n+1}$ is analytically continued along any path on D as a locally biholomorphic mapping. Let $\gamma : [0,1] \to D$ be a path on D. Then we take a piecewise line segment curve $\lambda : [0,1]$ and a continuous function $\Gamma : [0,1] \times [0,1] \to D$ such that

$$\begin{aligned} \gamma(0) &= \lambda(0), \quad \gamma(1) = \lambda(1) \\ \Gamma(0,\tau) &= \gamma(\tau), \quad \Gamma(1,\tau) = \lambda(\tau) \quad \text{for all } \tau \in [0,1] \end{aligned}$$

and the path $\Gamma(\cdot,\tau) : [0,1] \to D$ for each $\tau \in [0,1]$ is a piecewise line segment curve on D. Clearly, the mapping ϕ is analytically continued along the whole path $\gamma[0,1]$ so that the mapping ϕ is analytically continued along any path on D as a locally biholomorphic mapping.

Let G be the path space of D pointed by a point of $U \cap D$ mod homotopic relation so that G is a universal covering of D with a natural complex structure and a natural holomorphic covering map $\kappa : G \to D$. By the analytic continuation of the mapping $\phi : U \cap D \to B^{n+1}$ on D, the mapping $\varphi : B^{n+1} \to D$ has its natural proper locally biholomorphic lift $\psi : B^{n+1} \to G$. Since the mapping $\psi : B^{n+1} \to G$ is proper and locally biholomorphic, the mapping ψ is a finite covering of G. This completes the proof. □

Theorem 40. *Let D be a bounded domain in $\mathbb{C}^{n+1}$ with spherical differentiable boundary bD such that the fundamental group $\pi_1(D)$ is finite. Suppose that there is a biholomorphic mapping*

$$\phi \in H\left(U \cap D\right) \cap C^1\left(U \cap \overline{D}\right)$$

for a connected open neighborhood U of a point $p \in bD$ satisfying

$$\phi\left(U \cap bD\right) \subset bB^{n+1}.$$

Then D is necessarily simply connected and the mapping ϕ is analytically continued to a biholomorphic mapping from D onto B^{n+1}.

Proof. Let G be the path space of D pointed by a point of $U \cap D$ mod homotopic relation with a natural complex structure and a natural locally biholomorphic covering map $\kappa : G \to D$. Then, by Lemma 39, there is a locally biholomorphic finite covering lift $\psi : B^{n+1} \to G$ satisfying

$$\varphi = \kappa \circ \psi$$

where $\varphi : B^{n+1} \to D$ is an analytic continuation of the inverse mapping ϕ^{-1}.

Since the fundamental group $\pi_1\left(D\right)$ is finite, the mapping $\kappa : G \to D$ is a finite covering. Thus the mapping

$$\varphi = \kappa \circ \psi : B^{n+1} \to D$$

is a locally biholomorphic finite covering. Therefore, the analytic continuation of the inverse mapping ϕ^{-1} on the boundary bB^{n+1} yields finitely many germs at each point of bB^{n+1}. By Lemma 20, the Doggaebi variety L associated to the mapping ϕ is empty so that the mapping

$$\varphi = \phi^{-1} : bB^{n+1} \to bD$$

is also a locally biholomorphic finite covering. Thus the mapping

$$\varphi : \overline{B^{n+1}} \to \overline{D}$$

is a finite covering mapping. Then the fixed point property of the close ball $\overline{B^{n+1}}$ implies that the mapping $\varphi : \overline{B^{n+1}} \to \overline{D}$ is a simple cover. Otherwise, a nontrivial deck transformation of the closed ball $\overline{B^{n+1}}$ yields a continuous function on $\overline{B^{n+1}}$ with no fixed point. Therefore, the analytic continuation of the mapping $\phi : U \cap D \to B^{n+1}$ is analytically continued to a biholomorphic mapping

$$\varphi^{-1} : D \to B^{n+1}.$$

This completes the proof. □

3. Locally Biholomorphic Mappings

3.1. **Estimates of normalizing parameters.**

Lemma 41. *Let M be a nonspherical analytic real hypersurface in $\mathbb{C}^{n+1}$ and $Aut_p(M)$ be the isotropy subgroup at a point $p \in M$. Suppose that there is a real number $c \geq 1$ satisfying*

$$\sup_{\varphi \in Aut_p(M)} |U_\varphi| \leq c < \infty.$$

Then there is a real number $e \geq 1$ satisfying

$$|a_\phi| \leq e, \quad e^{-1} \leq |\rho_\phi| \leq e, \quad |r_\phi| \leq e$$

for every local automorphism $\phi \in Aut_p(M)$.

Proof. The real hypersurface $\mu_p(M)$ in normal form is expanded as follows:

$$v = \langle z, z\rangle + \sum_{k=4}^{\infty} F_k(z, \overline{z}, u)$$

where

$$F_k\left(\nu z, \nu\overline{z}, \nu^2 u\right) = \nu^k F_k(z, \overline{z}, u).$$

Since M is nonspherical, not all $F_k(z, \overline{z}, u)$ are zero. Then we make an estimate of the normalizing parameters $a_\phi, \rho_\phi, r_\phi$ for $\phi \in Aut_p(M)$(cf.[Pa3]). This completes the proof. □

Theorem 42. *Let M be a nonspherical strongly pseudoconvex analytic real hypersurface in a complex manifold. Then the local isotropy subgroup $Aut_p(M)$ is compact for every point $p \in M$.*

Proof. Because the situation is local, we may assume by taking a coordinate chart, if necessary, that the complex manifold is $\mathbb{C}^{n+1}$. Then the pseudoconvexity of M leads to

$$|U_\varphi| = 1 \quad \text{for} \quad \varphi \in Aut_p(M).$$

By Lemma 41, we obtain the desired result. This completes the proof. □

Lemma 43. *Let M, M' be nonspherical analytic real hypersurfaces in $\mathbb{C}^{n+1}$ and p, p' be points respectively of M, M' such that the two germs M at p and M' at p' are biholomorphically equivalent. Suppose that the isotropy subgroup $Aut_p(M)$ is compact. Then there is a real number $\delta_p > 0$ such that each germ of a biholomorphic mapping ϕ sending the germ M at p to the germ M' at p' is analytically continued to the open ball $B(p; \delta_p)$.*

Proof. We take a biholomorphic mapping ϕ on a connected open neighborhood U of the point $p \in M$ such that

$$\phi(U \cap M) \subset M'.$$

Then every germ of a biholomorphic mapping sending the germ M at p to the germ M' at p' is one of the following

$$\phi \circ \varphi \quad \text{for} \quad \varphi \in Aut_p(M).$$

Then the compactness of the group $Aut_p(M)$ leads to the desired result. This completes the proof. □

Lemma 44. *Let M be a nonspherical analytic real hypersurface in $\mathbb{C}^{n+1}$ such that the isotropy subgroup $Aut_p(M)$ is compact at every point $p \in M$. Then, for each compact subset $K \subset\subset M$, there is a real number $e \geq 1$ satisfying*

$$e^{-1} \leq |U_\phi| \leq e, \quad |a_\phi| \leq e, \quad e^{-1} \leq |\rho_\phi| \leq e, \quad |r_\phi| \leq e$$

for every point $p \in K$ and every local automorphism $\phi \in Aut_p(M)$.

Proof. The real hypersurface $\mu_p(M)$ in normal form is expanded as follows:

$$v = \langle z, z\rangle + \sum_{|I|,|J|\geq 2, k\geq 1} \lambda_{IJk}(p)\, z^I \overline{z}^J u^k.$$

Suppose that the normalization N_e, $e = (U, a, \rho, r) \in H$, transforms the real hypersurface $\mu_p(M)$ to a real hypersurfaces expanded as follows:

$$v = \langle z, z\rangle + \sum_{|I|,|J|\geq 2, k\geq 1} \eta_{IJk}(p; U, a, \rho, r)\, z^I \overline{z}^J u^k.$$

Note that the element $(U_\phi, a_\phi, \rho_\phi, r_\phi)$ for $\phi \in Aut_p(M)$ is characterized by the following equalities:

$$\lambda_{IJk}(p) = \eta_{IJk}(p; U, a, \rho, r), \quad |I|, |J| \geq 2, k \geq 1.$$

Then, since the algebraic subset of (U, a, ρ, r) is characterized by finitely many equalities, there is an integer K such that the following equalities

$$\lambda_{IJk}(p) = \eta_{IJk}(p; U, a, \rho, r), \quad |I|, |J|, k \leq K \tag{14}$$

characterize the element $(U_\phi, a_\phi, \rho_\phi, r_\phi)$ for $\phi \in Aut_p(M)$.

Since the isotropy subgroup $Aut_p(M)$ is compact, there is a real number $e_p \geq 1$ such that

$$e_p^{-1} \leq |U_\phi| \leq e_p, \quad |a_\phi| \leq e_p, \quad e_p^{-1} \leq |\rho_\phi| \leq e_p, \quad |r_\phi| \leq e_p$$

for every element $\phi \in Aut_p(M)$. Thus the algebraic set defined by the equalities in 14 is bounded. Note that the boundedness of the algebraic set is preserved on an open neighborhood of the point p so that there are a real number $\delta_p > 0$ and a real number $c_p \geq 1$ satisfying

$$c_p^{-1} \leq |U_\phi| \leq c_p, \quad |a_\phi| \leq c_p, \quad c_p^{-1} \leq |\rho_\phi| \leq c_p, \quad |r_\phi| \leq c_p$$

for every point $p \in B(p; \delta_p) \cap M$ and every local automorphism $\phi \in Aut_p(M)$.

Since the subset K is compact, there are finitely many points $p_1, \cdots, p_l$ such that

$$K \subset \bigcup_{k=1}^{l} B(p_k; \delta_{p_k}) \cap M.$$

Then we take

$$e = \max\{e_{p_k} : 1 \leq k \leq l\}.$$

This completes the proof. □

Lemma 45. *Let M be nonspherical analytic real hypersurfaces in $\mathbb{C}^{n+1}$ such that the isotropy subgroup $Aut_p(M)$ is compact at every point $p \in M$. Then, for each compact subset $K \subset\subset M$, there is a real number $\delta > 0$ such that each germ of a local automorphism $\phi \in Aut_p(M)$, $p \in K$, is analytically continued to the open ball $B(p; \delta)$.*

Proof. By the construction of the normalizing map μ_p, the real number δ_p depend only on the point $p \in M$. Hence there is a real number $\delta_p > 0$ for each $p \in M$ such that, for every $q \in B(p;\delta_p) \cap M$, the mapping μ_q is biholomorphic on $B(q;\delta_p)$ and the inverse μ_q^{-1} is biholomorphic on $B(0;\delta_p)$. Since K is a compact subset, there are finitely many points $p_1, \cdots, p_l$ such that

$$K \subset \bigcup_{k=1}^{l} B\left(p_k;\delta_{p_k}\right) \cap M.$$

Then we take

$$\delta_1 = \max\left\{\delta_{p_k} : 1 \leq k \leq l\right\}$$

so that μ_p is biholomorphic on $B(p;\delta_1)$ and μ_p^{-1} is biholomorphic on $B(0;\delta_1)$ for every $p \in K$.

By Lemma 44, there is a real number δ_2 such that every local automorphism $\varphi \in Aut_0(\mu_p(M))$ for every $p \in K$ is biholomorphically continued to the neighborhood $B(0;\delta_2)$. Then we take

$$\delta = \min\left\{\delta_1, \delta_2\right\}.$$

This completes the proof. □

Lemma 46. *Let M be a nonspherical analytic real hypersurface and $\gamma : [0,1] \rightarrow M$ be a chain-segment. Let M' be a nonspherical analytic real hypersurface in Moser-Vitushkin normal form(cf.* [Pa1]*). Suppose that there is a connected open neighborhood U of the point $\gamma(0)$ and a biholomorphic mapping ϕ on U such that*

$$\phi(U \cap M) \subset M'$$

and the image $\phi(U \cap \gamma[0,1])$ is on the straightened chain of M'. Then the mapping ϕ is analytically continued along the whole chain-segment $\gamma[0,1]$ as a local biholomorphic mapping.

Proof. Suppose that the assertion is not true. Then there is a real number λ, $0 < \lambda \leq 1$, such that the mapping ϕ is analytically continued along all the subpath $\gamma[0,\tau], \tau < \lambda$, but not the whole path $\gamma[0,\lambda]$, as a local biholomorphic mapping.

Since $\gamma[0,1]$ is a chain-segment, there is a biholomorphic mapping φ on an open neighborhood V of the point $\gamma(\lambda)$ such that the image $\varphi(V \cap \gamma[0,1])$ is on the straightened chain of M'. Then we take a point $p \in \varphi(V \cap \gamma[0,\lambda))$ and an open neighborhood W of the point p such that the mapping ϕ is analytically continued on W along the chain-segment $\gamma[0,1]$ and $W \subset V$ so that

$$\phi \circ \varphi^{-1}\left(\varphi(W) \cap M'\right) \subset M'$$

and the mapping $\phi \circ \varphi^{-1}$ maps the straightened chain $\varphi(W \cap \gamma[0,1])$ of M' onto the straightened chain of M'.

Note that the composition $\psi = \phi \circ \varphi^{-1}$ is necessarily analytically continued along the whole straightened chain of M'(cf. [Pa3]). Then, by abuse of notation, the composition $\psi \circ \varphi$ is the analytic continuation of the mapping ϕ on the point $\gamma(\lambda)$ along the chain-segment $\gamma[0,1]$. This is a contradiction. This completes the proof. □

3.2. **Analytic continuation on a real hypersurface.** We defined a canonical normalization ν_p of a nondegenerate analytic real hypersurface M at a point $p \in M$ to Moser-Vitushkin normal form(cf. [Pa1]) by the same way of the canonical normalization μ_p to Moser normal form.

Lemma 47. *Let M, M' be nonspherical analytic real hypersurfaces in complex manifolds such that M' is compact and the isotropy subgroup $Aut_p(M)$ is compact at every point $p \in M$. Let $\gamma : [0,1] \to M$ be a chain-segment and ϕ be a biholomorphic mapping on a connected open set U of the point $\gamma(0)$ such that*

$$\phi(U \cap M) \subset M'.$$

Then the mapping ϕ is analytically continued along the whole chain-segment $\gamma[0,1]$ as a local biholomorphic mapping.

Proof. Suppose that the assertion is not true. Then, without loss of generality, we may assume that the mapping ϕ is analytically continued along all the subpath $\gamma[0,\tau], \tau < 1$, but not the whole chain-segment $\gamma[0,1]$ as a local biholomorphic mapping. By the analytic continuation of the mapping ϕ, there is a path $\phi \circ \gamma : [0,1) \to M'$. Then we consider the following sequences

$$\begin{aligned} p_j &\equiv \gamma\left(1 - \frac{1}{j}\right), \quad j \in \mathbb{N}^+, \\ p'_j &\equiv \phi \circ \gamma\left(1 - \frac{1}{j}\right), \quad j \in \mathbb{N}^+. \end{aligned}$$

Since M' is compact, passing to a subsequence, say m_j, there is a point $p' \in M'$ such that $p'_{m_j} \to p'$ as $j \to \infty$ so that the closure

$$\overline{\left\{ p'_{m_j} \equiv \phi \circ \gamma\left(1 - \frac{1}{m_j}\right), \quad j \in \mathbb{N}^+ \right\}}$$

is a compact subset in a coordinate chart of the complex manifold.

Without loss of generality, we may assume that the chain $\gamma[0,1] \subset M$ in a coordinate chart of the complex manifold. By abuse of notation, we assume that M, M' are in $\mathbb{C}^{n+1}$. Let ν_p be the canonical normalization of a nondegenerate real hypersurface M at a point $p \in M$ to Moser-Vitushkin normal form.

Then the mapping ϕ yields the following germs of a biholomorphic mapping:

$$\varphi_j \equiv \nu_{p'_{m_j}} \circ \phi \circ \nu_{p_{m_j}}^{-1} : \nu_{p_{m_j}}(M) \to \nu_{p'_{m_j}}(M').$$

Since $\gamma : [0,1] \to M$ is a chain-segment, there is a sequence of local automorphisms

$$\sigma_j \in Aut_{p_{m_j}}(M)$$

such that the mapping $\nu_{p'_{m_j}} \circ \phi \circ \sigma_j$ sends the germ of the chain γ at the point p_{m_j} onto the straightened chain of the real hypersurface $\mu_{p'_{m_j}}(M')$ in Moser-Vitushkin normal form. By Lemma 46, the composition $\psi_j = \nu_{p'_{m_j}} \circ \phi \circ \sigma_j$ is analytically continued along the whole chain-segment $\gamma[0,1]$. Therefore, there is a real number $\delta > 0$ such that, by abuse of notation, the mapping ψ_j is biholomorphic on the open neighborhood $B\left(p_{m_j}; \delta\right)$.

By Lemma 45, the set $\left\{ \nu_{p_{m_j}} \circ \sigma_j \circ \nu_{p_{m_j}}^{-1} : j \in \mathbb{N}^+ \right\}$ is relatively compact.. Thus, by passing to a subsequence and shrinking $\delta > 0$, if necessary, we may assume that the composition $\tau_j = \nu_{p_{m_j}} \circ \sigma_j \circ \nu_{p_{m_j}}^{-1}$ and its converse τ_j^{-1} are biholomorphically continued on the open neighborhood $B(0;\delta)$.

Since $p_{m_j} \to \gamma(1)$ and $p'_{m_j} \to p'$, by passing to a subsequence and shrinking $\delta > 0$, if necessary, we may assume that the canonical normalizations $\nu_{p_{m_j}}, \nu_{p'_{m_j}}$ and their inverses $\nu_{p_{m_j}}^{-1}, \nu_{p'_{m_j}}^{-1}$ are biholomorphically continued respectively on the open neighborhoods

$$B\left(p_{m_j};\delta\right), \quad B\left(p'_{m_j};\delta\right), \quad B(0;\delta), \quad B(0;\delta).$$

Then, by abuse of notation, the composition

$$\chi_j = \nu_{p'_{m_j}}^{-1} \circ \psi_j \circ \nu_{p_{m_j}}^{-1} \circ \tau_j^{-1} \circ \nu_{p_{m_j}}$$

is biholomorphic on the open neighborhood $B\left(p_{m_j};\delta\right)$, if necessary, by shrinking $\delta > 0$.

Note that the mapping χ_j is a local biholomorphic continuation of the germ of the mapping ϕ at the point p_{m_j} for each $j \in \mathbb{N}^+$. Thus we take an integer K such that

$$\gamma(1) \in B\left(p_{m_K};\frac{\delta}{2}\right)$$

so that the mapping χ_K is an analytic continuation of the mapping ϕ on the point $\gamma(1)$ along the chain-segment $\gamma[0,1]$. This contradiction completes the proof. □

Lemma 48. *Let M, M' be nonspherical analytic real hypersurfaces in complex manifolds such that M' is compact and the isotropy subgroup $Aut_p(M)$ is compact at every point $p \in M$. Suppose that there is a biholomorphic mapping ϕ on a connected open set U of a point $p \in M$ such that*

$$\phi(U \cap M) \subset M'.$$

Then the biholomorphic mapping ϕ is analytically continued along any path on M as a local biholomorphic mapping.

Proof. For each point $p \in M$, we make a biholomorphically equivalent deformation of the real hypersurfaces $\mu_p(M)$ in normal form continuously to a real hyperquadric by using the scaling mapping

$$\begin{cases} z^* = \lambda z \\ w^* = \lambda^2 w \end{cases}, \quad \lambda \in \mathbb{R}^+.$$

Because the chain is characterized by an order differential equation, the continuous family of chains on the real hyperquadric is continuously deformed by the parameter λ on a real hypersurface biholomorphic to M near the point p(cf. [Pa3]).

Let $\gamma : [0,1] \to M$ be a path on M such that $\gamma(0) \in U \cap M$. Then, for each $\tau \in [0,1]$, there is a real number $\varepsilon_\tau > 0$, a point $p_\tau \in M$ and a continuous function

$$\Gamma_\tau : [0,1] \times ([0,1] \cap (\tau - \varepsilon_\tau, \tau + \varepsilon_\tau)) \to M$$

such that

1. $\Gamma_\tau(\cdot,\sigma) : [0,1] \to M$ is a chain-segment for each $\sigma \in [0,1] \cap (\tau - \varepsilon_\tau, \tau + \varepsilon_\tau)$,
2. $\Gamma_\tau(0,\sigma) = \gamma(\sigma)$ for each $\sigma \in [0,1] \cap (\tau - \varepsilon_\tau, \tau + \varepsilon_\tau)$,
3. $\Gamma_\tau(1,\sigma) = p_\tau$ for all $\sigma \in [0,1] \cap (\tau - \varepsilon_\tau, \tau + \varepsilon_\tau)$.

Note that the family $\{[0,1]\cap(\tau-\varepsilon_\tau,\tau+\varepsilon_\tau):\tau\in[0,1]\}$ is an open covering of the compact set $[0,1]$. Thus there is a finite subcover

$$\left\{[0,1]\cap(\tau_j-\varepsilon_{\tau_j},\tau_j+\varepsilon_{\tau_j}):\tau_j\in[0,1],\quad j=1,\cdots,m\right\}.$$

Then, by Lemma 47, the biholomorphic mapping ϕ is analytically continued along the whole path $\gamma[0,1]$ as a local biholomorphic mapping. This completes the proof. □

3.3. **Holomorphic mapping on the boundary.**

Lemma 49. *Let Q be a real hyperquadric defined by*

$$v=\langle z,z\rangle\equiv z^1\overline{z}^1+\cdots+z^n\overline{z}^n$$

and ϕ be a polynomial mapping as follows:

$$\phi:\begin{cases} z^*=f(z,w)\\ w^*=g(z,w)\end{cases}$$

where, for $m\geq 2$,

$$\begin{aligned} f(\mu z,\mu^2w) &= \mu^m f(z,w)\\ g(\mu z,\mu^2w) &= \mu^{2m}g(z,w). \end{aligned}\tag{15}$$

Suppose that

$$\phi(Q)\subset Q.$$

Then $\phi\equiv 0$.

Proof. The mapping $\phi=(f,g)$ yields the identity

$$\Im g(z,u+i\langle z,z\rangle)=\langle f(z,u+i\langle z,z\rangle),f(z,u+i\langle z,z\rangle)\rangle.\tag{16}$$

Suppose that m is even, i.e., $m=2k\geq 2$. We may consider $z,\overline{z},u$ as independent variables in the identity 16. Taking $\overline{z}=0$ in the equality 16 yields

$$g(z,u)-\overline{g}(0,u)=2i\langle f(z,u),f(0,u)\rangle.$$

Thus we can put

$$g(z,u)=2i\langle f(z,u),f(0,u)\rangle+\Re g(0,u)-i\langle f(0,u),f(0,u)\rangle.\tag{17}$$

Note that

$$f(0,u)=f(0,1)u^{\frac{m}{2}},\quad g(0,u)=g(0,1)u^m.$$

Then the identity 16 yields

$$\begin{aligned} &2\Re\left\{\langle f(z,u+i\langle z,z\rangle),f(0,1)\rangle(u+i\langle z,z\rangle)^{\frac{m}{2}}\right\}\\ &+\Re g(0,1)\Im(u+i\langle z,z\rangle)^m\\ &-\langle f(0,1),f(0,1)\rangle\Re(u+i\langle z,z\rangle)^m\\ =\ &\langle f(z,u+i\langle z,z\rangle),f(z,u+i\langle z,z\rangle)\rangle. \end{aligned}$$

Let $p(z,\overline{z})$ be a polynomial of the variables $z,\overline{z}$ satisfying

$$p(\mu z,\nu\overline{z})=\mu^l\nu^m p(z,\overline{z}).$$

Then the polynomial $p(z,\overline{z})$ is said to be of type (l,m). Collecting the terms of type $(l,1)$, $l=0,1,\cdots,m-1$, yields

$$miu^{-1}\langle z,z\rangle\{\langle f(z,u),f(0,u)\rangle-\langle f(0,u),f(0,u)\rangle+\Re g(0,u)\}=0.$$

Hence we obtain

$$\langle f(z,u), f(0,u)\rangle - \langle f(0,u), f(0,u)\rangle + \Re g(0,u) = 0,$$

so that

$$\begin{array}{rcl}\langle f(z,u), f(0,u)\rangle &=& \langle f(0,u), f(0,u)\rangle\\ \Re g(0,u) &=& 0.\end{array}$$

Thus, from the identity 17, we obtain

$$g(z,u) = i\langle f(0,u), f(0,u)\rangle$$

by which the identity 16 yields

$$\langle f(0,1), f(0,1)\rangle \Re(u+i\langle z,z\rangle)^m = \langle f(z,u+i\langle z,z\rangle), f(z,u+i\langle z,z\rangle)\rangle.$$

Note that

$$\langle f(z,u+i\langle z,z\rangle), f(z,u+i\langle z,z\rangle)\rangle \geq 0,$$

but

$$\Re(u+i\langle z,z\rangle)^m = u^m - \frac{m(m-1)}{2}u^{m-2}\langle z,z\rangle^2 + \cdots .$$

Thus collecting terms of type $(2,2)$ yields

$$\begin{array}{rl} & -\dfrac{m(m-1)}{2}\langle f(0,1), f(0,1)\rangle u^{m-2}\langle z,z\rangle^2\\ = & \left|\displaystyle\sum_{\alpha,\beta}\frac{z^\alpha z^\beta}{2}\left(\frac{\partial^2 f}{\partial z^\alpha \partial z^\beta}\right)(0,u)\right|^2 + \dfrac{mu^{-2}}{2}\langle z,z\rangle^2\langle f(0,u), f(0,u)\rangle\\ \geq & 0.\end{array}$$

Since $m \geq 2$, we obtain

$$\langle f(0,1), f(0,1)\rangle = 0$$

which yields

$$\langle f(z,u+i\langle z,z\rangle), f(z,u+i\langle z,z\rangle)\rangle = 0,$$

i.e.,

$$f(z,w) = g(z,w) = 0.$$

Suppose that m is odd, i.e., $m = 2k+1 \geq 3$. We may consider $z, \overline{z}, u$ as independent variables in the identity 16. Then, by the condition 15, taking $\overline{z} = 0$ in the equality 16 yields

$$g(z,u) = \overline{g}(0,u)$$

since

$$f(0,u) = 0.$$

Thus we can put

$$g(z,u) = g(0,u), \quad g(0,1) \in \mathbb{R}.$$

Then the identity 16 yields

$$g(0,1)\Im(u+i\langle z,z\rangle)^m = \langle f(z,u+i\langle z,z\rangle), f(z,u+i\langle z,z\rangle)\rangle. \tag{18}$$

Note that

$$\langle f(z,u+i\langle z,z\rangle), f(z,u+i\langle z,z\rangle)\rangle \geq 0,$$

but

$$\begin{aligned}\Im(u+i\langle z,z\rangle)^m &= mu^{m-1}\langle z,z\rangle \\ &\quad -\frac{m(m-1)(m-2)}{2}u^{m-3}\langle z,z\rangle^3+\cdots .\end{aligned}$$

Thus collecting terms of type $(1,1)$ yields

$$\begin{aligned}&mg(0,1)u^{m-1}\langle z,z\rangle \\ =& \left|\sum_{\alpha} z^{\alpha}\left(\frac{\partial f}{\partial z^{\alpha}}\right)(0,u)\right|^2 \geq 0.\end{aligned}$$

Collecting terms of type $(3,3)$ yields

$$\begin{aligned}&-\frac{m(m-1)(m-2)}{2}g(0,1)u^{m-3}\langle z,z\rangle^3 \\ =& \left|\sum_{\alpha,\beta,\gamma}\frac{z^{\alpha}z^{\beta}z^{\gamma}}{6}\left(\frac{\partial^3 f}{\partial z^{\alpha}\partial z^{\beta}\partial z^{\gamma}}\right)(0,u)\right|^2 \\ &+\frac{m-1}{2}\left|\sum_{\alpha} z^{\alpha}\left(\frac{\partial f}{\partial z^{\alpha}}\right)(0,u)\right|^2 u^{-2}\langle z,z\rangle^2 \\ \geq& \ 0.\end{aligned}$$

Since $m \geq 3$, we obtain

$$g(0,1)\langle z,z\rangle = 0$$

by which the identity 18 yields

$$\langle f(z,u+i\langle z,z\rangle), f(z,u+i\langle z,z\rangle)\rangle = 0,$$

i.e.,

$$f(z,w) = g(z,w) = 0.$$

This completes the proof. □

Lemma 50. *Let M, M' be strongly pseudoconvex analytic real hypersurfaces in $\mathbb{C}^{n+1}$. Let ϕ be a holomorphic mapping on an open neighborhood U of a point $p \in M$ such that*

$$\phi(U \cap M) \subset M'.$$

Then the mapping ϕ is either a constant mapping or a biholomorphic mapping on U, if necessary, shrinking U.

Proof. We take $q = \phi(p)$ so that

$$\varphi \equiv \mu_q \circ \phi \circ \mu_p^{-1} : \mu_p(M) \to \mu_q(M').$$

Then the mapping φ is a holomorphic mapping on an open neighborhood V of the origin satisfying

$$\varphi(\mu_p(M) \cap V) \subset \mu_q(M').$$

The mapping $\varphi=(f,g)$ in $\mathbb{C}^n\times\mathbb{C}$ is decomposed as follows:

$$f(z,w)=\sum_{k=1}^{\infty}f_k(z,w),\quad g(z,w)=\sum_{k=1}^{\infty}g_k(z,w),$$

where

$$f_m(\mu z,\mu^2 w)=\mu^m f_m(z,w),\quad g_m(\mu z,\mu^2 w)=\mu^m g_m(z,w).$$

We assume that $\mu_p\left(M\right),\mu_q\left(M'\right)$ are defined respectively by the equations

$$v=F(z,\overline{z},u),\quad v=\langle z,z\rangle+F^*(z,\overline{z},u)$$

where

$$F(z,\overline{z},u)=\langle z,z\rangle+O\left(|z|^4\right),\quad F^*(z,\overline{z},u)=O\left(|z|^4\right).$$

Then we obtain the following identity near the origin

$$\begin{aligned} & \Im g(z,u+iF(z,\overline{z},u)) \\ = & \langle f(z,u+iF(z,\overline{z},u)),f(z,u+iF(z,\overline{z},u))\rangle \\ & +F^*(f(z,u+iF(z,\overline{z},u)),\overline{f(z,u+iF(z,\overline{z},u))},\Re g(z,u+iF(z,\overline{z},u))). \end{aligned} \tag{19}$$

Then, up to weight 2, we obtain

$$\Im g_1(z,0)=0$$
$$\Im g_2(z,u+i\langle z,z\rangle)=\langle f_1(z,0),f_1(z,0)\rangle$$

which yields

$$\langle f_1(z,0),f_1(z,0)\rangle=\langle z,z\rangle\Re g_2(0,1)$$

and

$$\begin{aligned} g_1(z,0) &= 0 \\ g_2(z,w) &= g_2(0,1)w,\quad g_2(0,1)\in\mathbb{R}. \end{aligned}$$

Note that $g_2(0,1)\neq 0$ if and only if the mapping φ is a biholomorphic mapping at the origin.

Suppose that the assertion is not true. Then we have

$$f_1(z,0)=g_1(z,0)=g_2(z,w)=0.$$

As inductive hypothesis, suppose that, for $m\geq 2$,

$$\begin{aligned} f_l(z,w) &= 0,\quad l=1,\cdots,m-1, \\ g_l(z,w) &= 0,\quad l=1,\cdots,2m-2. \end{aligned} \tag{20}$$

By the condition

$$F(z,\overline{z},u)=\langle z,z\rangle+O\left(|z|^4\right),\quad F^*(z,\overline{z},u)=o\left(|z|^4\right),$$

the identity 19 yields

$$\Im g_{2m-1}(z,u+i\langle z,z\rangle)=0$$

Here we may consider $z,\overline{z},u$ as independent variables so that taking $\overline{z}=0$ yields

$$g_{2m-1}(z,u+i\langle z,z\rangle)=\overline{g_{2m-1}}(0,u-i\langle z,z\rangle)=0.$$

Hence the hypothesis 20 necessarily comes to

$$\begin{aligned} f_l(z,w) &= 0, \quad l = 1,\cdots,m-1, \\ g_l(z,w) &= 0, \quad l = 1,\cdots,2m-1. \end{aligned}$$

Then the identity 19 yields

$$\Im g_{2m}(z, u + i\langle z,z\rangle) = \langle f_m(z, u + i\langle z,z\rangle), f_m(z, u + i\langle z,z\rangle)\rangle.$$

Note that the polynomial mapping $\varphi_m \equiv (f_m, g_{2m})$, $m \geq 2$, satisfies

$$\varphi_m(Q) \subset Q.$$

By Lemma 49, $\varphi_m \equiv 0$ so that

$$\begin{aligned} f_l(z,w) &= 0, \quad l = 1,\cdots,m, \\ g_l(z,w) &= 0, \quad l = 1,\cdots,2m. \end{aligned}$$

This completes the induction so that $\varphi \equiv 0$, i.e., the mapping φ is a constant mapping. This completes the proof. □

3.4. **Proper holomorphic mappings.**

Lemma 51. *Let D, D' be strongly pseudoconvex bounded domains with nonspherical real analytic boundaries bD, bD' such that the boundaries bD, bD' are both simply connected. Suppose that there is a biholomorphic mapping ϕ on a connected open neighborhood U of a point $p \in bD$ such that*

$$\phi(U \cap bD) \subset bD'.$$

Then the mapping ϕ is analytically continued to a biholomorphic mapping from D onto D'.

Proof. Since bD, bD' are simply connected, by Lemma 48, the mappings ϕ, ϕ^{-1} are both analytically continued, by abuse of notation, respectively to a biholomorphic mapping

$$\phi : D \to D$$

and

$$\phi^{-1} : D' \to D.$$

Then, by the identity theorem, the mapping ϕ is biholomorphic. This completes the proof. □

Lemma 52. *Let D, D' be strongly pseudoconvex bounded domains with nonspherical real analytic boundaries bD, bD' such the boundary bD' is simply connected and the closed set $\overline{D'}$ satisfies the fixed point property. Suppose that there is a biholomorphic mapping ϕ on a connected open neighborhood U of a point $p \in bD$ such that*

$$\phi(U \cap bD) \subset bD'.$$

Then the mapping ϕ is analytically continued to a biholomorphic mapping from D onto D'.

Proof. Since bD' is simply connected, by Lemma 48, the inverse mapping ϕ^{-1} is analytically continued to a locally biholomorphic proper mapping

$$\varphi : D' \to D.$$

Note that the mappings $\varphi : D' \to D$ and $\varphi : bD' \to bD$ are both covering maps. We claim that the covering is simple. Otherwise, there is a nontrivial deck transformation of $\overline{D'}$ yields a continuous self mapping of $\overline{D'}$ with no fixed point. This is a contradiction. Therefore, the mapping $\varphi : D' \to D$ is biholomorphic. This completes the proof. □

Lemma 53. *Let D, D' be strongly pseudoconvex bounded domains with nonspherical real analytic boundaries bD, bD' such that the domain D and the boundary bD' are both simply connected. Suppose that there is a biholomorphic mapping ϕ on a connected open neighborhood U of a point $p \in bD$ such that*

$$\phi\,(U \cap bD) \subset bD'.$$

Then the mapping ϕ is analytically continued to a biholomorphic mapping from D onto D'.

Proof. Since bD' is simply connected, by Lemma 48, the inverse mapping ϕ^{-1} is analytically continued to a locally biholomorphic proper mapping

$$\varphi : D' \to D.$$

Note that the mapping $\varphi : D' \to D$ is a covering map. Since D is simply connected, the covering is simple. Therefore, the mapping $\varphi : D' \to D$ is biholomorphic. This completes the proof. □

Theorem 54. *Let D, D' be strongly convex bounded domains with real analytic boundaries bD, bD'. Suppose that there is a biholomorphic mapping ϕ on a connected open neighborhood U of a point $p \in bD$ such that*

$$\phi\,(U \cap bD) \subset bD'.$$

Then the mapping ϕ is analytically continued to a biholomorphic mapping from D onto D'.

Proof. Note that a strongly convex bounded domain is homeomorphic to an open ball B^{n+1}. Suppose that the boundaries bD, bD' are spherical. Then we take a biholomorphic mapping φ on U, if necessary, shrinking U, such that

$$\varphi\,(U \cap bD) \subset bB^{n+1}.$$

Then, by Lemma 19, the mapping φ and the composition $\psi \equiv \varphi \circ \phi^{-1}$ are analytically continued, by abuse of notation, to biholomorphic mappings as follows:

$$\varphi : D \to B^{n+1} \quad \text{and} \quad \psi : D' \to B^{n+1}.$$

Thus the composition $\psi^{-1} \circ \varphi : D \to D'$ is a biholomorphic mapping and the analytic continuation of the mapping ϕ.

Suppose that the boundaries bD, bD' are nonspherical. Then, by Lemma 53, the mapping ϕ is analytically continued to a biholomorphic mapping from D onto D'. This completes the proof. □

Lemma 55. *Let D, D' be strongly pseudoconvex bounded domains with real analytic boundaries bD, bD'. Suppose that there is a proper holomorphic mapping $\phi : D \to D'$. Then there is an open neighborhood U of a point $p \in bD$ such that the mapping ϕ is analytically continued on U and*

$$\phi (U \cap bD) \subset bD'.$$

Proof. We may apply the boundary regularity of Lemma 16 so that the mapping $\varphi : D \to D'$ is holomorphic on an open neighborhood of $\overline{D}$. This completes the proof. $\square$

Lemma 56. *Let $\phi : B^{n+1} \to B^{n+1}$ be a proper holomorphic mapping. Then $\phi \in Aut \left(B^{n+1}\right)$.*

Proof. By Lemma 55, there is an open neighborhood U of a point $p \in bB^{n+1}$ such that ϕ is analytically continued on U and

$$\phi \left(U \cap bB^{n+1}\right) \subset bB^{n+1}.$$

By Lemma 17, $\phi \in Aut \left(B^{n+1}\right)$. This completes the proof. $\square$

Lemma 57. *Let D, D' be strongly pseudoconvex bounded domains in $\mathbb{C}^{n+1}$ with real analytic boundaries bD, bD'. Suppose that there is a proper holomorphic mapping $\phi : D \to D'$. Then the mapping $\phi : D \to D'$ is a locally biholomorphic mapping so that $\phi : \overline{D} \to \overline{D'}$ is a covering map.*

Proof. By Lemma 55, there is a point $p \in bD$ and an open neighborhood U of the point p such that the mapping ϕ is analytically continued to U and

$$\phi (U \cap bD) \subset bD'.$$

By Lemma 50, the mapping ϕ is biholomorphic on U, if necessary, shrinking U.

Suppose that the boundaries bD, bD' are spherical. Then, by Lemma 30, the mapping $\phi : D \to D'$ is a locally biholomorphic mapping.

Suppose that the boundary bD, bD' are nonspherical. Then, by Lemma 48, the mapping $\phi : D \to D'$ is analytically continued on an open neighborhood of the boundary bD to be locally biholomorphic. Thus the mapping ϕ is a locally biholomorphic mapping. This completes the proof. $\square$

Theorem 58. *Let D be a strongly pseudoconvex bounded domain with real analytic boundary bD. Suppose that there is a proper holomorphic self mapping $\phi : D \to D$. Then ϕ is a biholomorphic automorphism of D.*

Proof. By Lemma 57, the mapping $\phi : \overline{D} \to \overline{D}$ is a self covering map. We claim that $\phi : D \to D$ is a simple covering. Otherwise, there would be a integer $m > 1$ such that, for every $p \in \overline{D}$, m is the order of the set

$$\left\{q \in \overline{D} : \phi (q) = p\right\}.$$

Then we define the k times composition ϕ^k of the mapping ϕ as follows

$$\phi^k \equiv \underbrace{\phi \circ \cdots \circ \phi}_{k} : \overline{D} \to \overline{D}.$$

Note that the inverse image of each point $p \in \overline{D}$ under the mapping $\phi^k : \overline{D} \to \overline{D}$, $k \in \mathbb{N}^+$, is a set of order m^k.

We claim that, for a given real number $\delta > 0$, there is a point $q \in bD$ and a compact subset $K \subset\subset D$ and a subsequence ϕ^{m_k} such that

$$\phi^{-m_k}(B(q;\delta) \cap D) \cap K \neq \emptyset \quad \text{for all } k \in \mathbb{N}^+.$$

Otherwise, for every compact subset $K \subset\subset D$, there is an integer l such that

$$\phi^{-k}(D \backslash K) \subset D \backslash K \quad \text{for all } k \geq l.$$

Since $\phi^k : D \to D$ is a finite covering map, we have

$$D = \phi^{-k}(K \cup D \backslash K) = \phi^{-k}(K) \cup \phi^{-k}(D \backslash K)$$

so that

$$K \subset \phi^{-k}(K) \quad \text{for all } k \geq l.$$

Note that the inverse image of $\phi^{-k}(K)$ is a union of m^k disconnected compact subsets. We may assume that K is connected so that K is in a compact subset of the union $\phi^{-k}(K)$. This is impossible. Thus, with such a point $q \in bD$, we take an accumulation point $q' \in D'$ of the set

$$\lim_{k \to \infty} \phi^{-m_k}(\{q\}) = \lim_{k \to \infty} \{p \in bD : \phi^{m_k}(p) = q\}.$$

Suppose that bD is spherical. Then there are open neighborhoods U, U' respectively of q, q' and biholomorphic mappings ψ, ψ' respectively on U, U' such that

$$\begin{aligned} \psi(U \cap bD) &\subset bB^{n+1} \\ \psi'(U' \cap bD) &\subset bB^{n+1}. \end{aligned}$$

By Lemma 17, the compositions

$$\varphi_k \equiv \psi' \circ \phi^{-k} \circ \psi^{-1} : \psi(U \cap D) \to \psi'(U' \cap D)$$

is analytically continued to, by abuse of notation, automorphisms $\varphi_k \in Aut\left(B^{n+1}\right)$. By the construction, a subsequence φ_{m_k} must converges to the point $\psi'(q')$ uniformly on every compact subset of B^{n+1}. Thus there is a sequence of real numbers $\delta_k \searrow 0$ such that

$$\phi^{m_k} : B(q';\delta_k) \cap D \to U \cap D$$

and

$$\phi^{m_k} = \psi^{-1} \circ \varphi_{m_k}^{-1} \circ \psi' \quad \text{on } B(q';\delta_k) \cap D.$$

Then we take a point $p \in U \cap D$ such that there is a sequence q_k satisfying

$$q_k \in \phi^{-m_k}(\{p\}) \cap B(q';\delta_k)$$

and, for a compact subset $K \subset\subset D$, there is a sequence p_k satisfying

$$p_k \in \phi^{-m_k}(\{p\}) \cap K.$$

Therefore, there is a sequence of deck transformations $\phi_k \in Aut(D)$ of the covering map $\phi^{m_k} : D \to D$ such that

$$\phi_k(p_k) = q_k \to q' \in bD.$$

Since $p_k \in K$, by Lemma 15, there is a biholomorphic mapping

$$\sigma : D \to B^{n+1}.$$

Then the composition

$$\iota \equiv \sigma \circ \phi \circ \sigma^{-1} : B^{n+1} \to B^{n+1}$$

would be a proper self mapping, but not an automorphism of the unit ball B^{n+1}. This is a contradiction to Lemma 56 that every proper self mapping of the unit ball B^{n+1} is an automorphism of the unit ball B^{n+1}.

Suppose that bD is nonspherical. We take a sequence $p_k \in bD$ such that

$$p_k \in \phi^{-k}\left(\{q\}\right) \subset bD$$

and

$$\left|p_k - q'\right| = \min\left\{\left|p - q'\right| : p \in \phi^{-k}\left(\{q\}\right)\right\}.$$

Let μ_{p_k} be the canonical normalizing mapping at the point $p_k \in bD$. Since $p_k \to q'$, there is an open neighborhood W of the point q' such that the mapping μ_{p_k} is analytically continued on W. Then we define a function $\varepsilon_k(p)$ for a sufficiently large k and a point $p \in W \cap bD$ such that

$$\varepsilon_k(p) \equiv \sum_{j=1}^{n} \left|z_j \circ \mu_{p_k}(p)\right|^2 + \left|w \circ \mu_{p_k}(p)\right|$$

where z_j, $j = 1, \cdots, n$, w are the coordinate functions of $\mathbb{C}^{n+1}$ such that z_j, $j = 1, \cdots, n$, are for the complex tangent hyperplane and w are for the complex line normal to the complex tangent hyperplane of $\mu_{p_k}(W \cap bD)$ at the origin. Then we take a sequence q_k such that

$$q_k \in \phi^{-k}\left(\{q\}\right) \subset bD$$

and

$$\varepsilon_k(q_k) = \min\left\{\varepsilon_k(p) : p \in \phi^{-k}\left(\{q\}\right)\right\}.$$

Let π_k be a complex line containing p_k and q_k. Then we take a subsequence π_{m_k} so that π_{m_k} converges to a complex line $\pi_{q'} \subset T_{q'}\mathbb{C}^{n+1}$. In other words, the a subsequence q_{m_k} converges to the point q' to a direction. Then we obtain

$$\varphi_k \equiv \mu_{p_k} \circ \phi^{-k} \circ \mu_q^{-1} : \mu_q(bD) \to \mu_{p_k}(bD).$$

Then we define a biholomorphic mapping

$$\sigma_k : \begin{cases} z^* = \sqrt{\varepsilon_{m_k}}z \\ w^* = \varepsilon_{m_k}w \end{cases}$$

where

$$\varepsilon_{m_k} = \varepsilon_{m_k}(q_{m_k}).$$

Then we take the composition

$$\kappa_k \equiv \sigma_k^{-1} \circ \varphi_{m_k} : \mu_q(bD) \to \sigma_k^{-1} \circ \mu_{p_{m_k}}(bD).$$

Note that there is a real number $\delta > 0$ such that the germs of real hypersurfaces $\mu_q(bD)$ and $\sigma_k^{-1} \circ \mu_{p_{m_k}}(bD)$ are analytically continued to the open neighborhood $B(0;\delta)$. Since the boundary bD is nonspherical, by Lemma 45, the mapping κ_k is biholomorphic on $B(0;\delta)$, if necessary, shrinking $\delta > 0$. Further, by the construction, the limit of the sequence κ_k cannot be a constant mapping on $B(0;\delta)$ so that, by Theorem 50, the limit of the sequence κ_k would be biholomorphic on $B(0;\delta)$.

By the way, the sequence of real hypersurfaces $\sigma_k^{-1} \circ \mu_{p_{m_k}}(bD)$ converges to a real hyperquadric uniformly on an open neighborhood of the origin. Therefore, the germ of the boundary bD at the point q' is spherical so that the boundary bD is spherical(cf. [Pa3]). This is a contradiction to the fact that the boundary bD is nonspherical. This completes the proof. □

3.5. **Locally realizable CR manifolds.** Let M be a CR manifold of CR dimension n and CR codimension 1 with a CR structure (D, I) where D is $2n$ dimensional smooth subbundle of the tangent bundle TM and I is an automorphism on D such that

$$I^2 V = -V \quad \text{for} \quad V \in \Gamma D.$$

For each point $p \in M$, there is a local coordinate chart (U, φ) such that

$$\varphi(U) \subset \mathbb{R}^{2n+1}.$$

Then M shall be called locally realizable CR manifold if there is an open neighborhood U of each point $p \in M$ and CR functions $f_1, \cdots, f_{n+1}$ on $\varphi(U)$ satisfying

$$df_1 \wedge \cdots \wedge df_{n+1} \neq 0.$$

Let z_j, $j = 1, \cdots, n+1$, be the coordinate functions of $\mathbb{C}^{n+1}$. Then we obtain a local embedding σ of $U \subset M$ into $\mathbb{C}^{n+1}$ such that

$$f_j \equiv z_j \circ \sigma \circ \varphi^{-1}.$$

The CR manifold M shall be called a locally analytically realizable CR manifold when $\sigma(U)$ is real analytic by a local embedding σ on an open neighborhood U of each point $p \in M$. Note that M is either spherical or nonspherical(cf. [Pa3]).

Lemma 59. *Let M be a connected locally analytically realizable CR manifold. Suppose that there is a nontrivial CR mapping φ on an open neighborhood U of a point $p \in M$ such that*

$$\varphi(U) \subset bB^{n+1}.$$

Then the mapping φ is CR continued along any path on M as a locally CR diffeomorphic mapping.

Proof. Since M is connected, M is necessarily spherical(cf. [Pa3]). Suppose that the assertion is not true. Then there is a path $\gamma : [0, 1] \to M$ with $\gamma(0) = p$ such that the mapping φ is CR continued along all subpath $\gamma[0, \tau]$ with $\tau < 1$, but not the whole path $\gamma[0, 1]$. Then we take a CR embedding $\sigma(V) \subset \mathbb{C}^{n+1}$ of an open neighborhood V of the point $\gamma(1)$ such that $\sigma(V)$ is a spherical analytic real hypersurface and

$$\varphi \circ \sigma^{-1} : \sigma(V) \to bB^{n+1}$$

is a locally CR diffeomorphism. Without loss of generality, we may assume that $\gamma[0, 1] \subset V$. Hence there is an open neighborhood W of the point $\sigma(\gamma(0))$ and a biholomorphic mapping ϕ on W such that

$$\phi = \varphi \circ \sigma^{-1} \quad \text{on } \sigma(V) \cap W.$$

Then the mapping ϕ is analytically continued along the whole path $\sigma(\gamma[0, 1])$. From, by abuse of notation, the mapping ϕ at an open neighborhood W' of the point $\sigma(\gamma(1))$ and a CR embedding σ of an open neighborhood V' of the point $\gamma(1)$, we obtain the CR mapping

$$\phi \circ \sigma : \sigma^{-1}(W' \cap \sigma(V')) \to bB^{n+1}$$

which is a CR continuation of the CR mapping φ. This completes the proof. ☐

Theorem 60. *Let M be a connected locally analytically realizable CR manifold. Suppose that M is compact and there is a nontrivial CR mapping φ on an open neighborhood U of a point $p \in M$ such that*

$$\varphi\left(U\right) \subset bB^{n+1}.$$

Then there is a finite subset $L \subset bB^{n+1}$ such that the inverse CR mapping φ^{-1} is CR continued along any path on $bB^{n+1} \backslash L$ as a locally CR diffeomorphic mapping.

Proof. Since M is connected, M is necessarily spherical(cf. [Pa3]). Let $q_j \in bB^{n+1}$ be a sequence converging to a point $q \in bB^{n+1}$ and ϕ_j be a sequence of the CR continuation of the inverse mapping φ^{-1} at the point q_j. We set

$$q_j' = \phi_j\left(q_j\right) \in M.$$

Since M is compact, there is a subsequence q_{m_j}' and a point $q' \in M$ such that

$$q_{m_j}' \to q'.$$

Then we take a CR embedding σ of an open neighborhood V of the point $q' \in M$ such that $\sigma\left(V\right) \subset \mathbb{C}^{n+1}$ is a spherical analytic real hypersurface. Then we apply the same argument as in the previous sections so that the singular locus L is a finite subset of bB^{n+1} and the inverse CR mapping φ^{-1} is CR continued along any path on $bB^{n+1} \backslash L$. This completes the proof. $\square$

Lemma 61. *Let M, M' be connected locally analytically realizable CR manifolds with positive definite Levi form. Suppose that M, M' are nonspherical and M' is compact, and there is a nontrivial CR mapping φ on an open neighborhood U of a point $p \in M$ such that*

$$\varphi\left(U\right) \subset M'.$$

Then the mapping φ is CR continued along any path on M as a locally CR diffeomorphic mapping.

Proof. Since M, M' are locally analytically realizable and the Levi forms of M, M' are positive definite, the isotropy subgroups $Aut_p\left(M\right), Aut_{p'}\left(M'\right)$ are compact for every point $p \in M, p' \in M'$.

Let $q_j \in M$ be a sequence converging to a point $q \in M$ and φ_j be a sequence of the CR continuation of the mapping φ at the point q_j. We set

$$q_j' = \varphi_j\left(q_j\right) \in M'.$$

Since M' is compact, there is a subsequence q_{m_j}' and a point $q' \in M'$ such that

$$q_{m_j}' \to q'.$$

Then we take CR embeddings σ, σ' respectively of open neighborhoods V, V' respectively of the points $q \in M, q' \in M'$ such that $\sigma\left(V\right), \sigma'\left(V'\right) \subset \mathbb{C}^{n+1}$ are nonspherical analytic real hypersurface. Then we apply the same argument as in the previous subsection. This completes the proof. $\square$

Theorem 62. *Let M, M' be connected locally analytically realizable CR manifolds with positive definite Levi form. Suppose that M, M' are compact and nonspherical, and there is a nontrivial CR mapping φ on an open neighborhood U of a point $p \in M$ such that*

$$\varphi\left(U\right) \subset M'.$$

Then the maximal CR extension of the mapping φ is a CR equivalence between the natural universal covering spaces of M, M' to be the pointed path spaces respectively of M, M' mod homotopic relation.

Proof. By Lemma 61, the mapping φ is CR continued along any path on M as a locally CR diffeomorphic mapping. Note that M is compact and

$$\varphi^{-1}\left(\varphi\left(U\right)\right) \subset M$$

so that we apply Lemma 61 to the inverse mapping φ^{-1}. Thus the inverse mapping φ^{-1} is CR continued along any path on M' as a locally CR diffeomorphic mapping. Thus the CR continuation of the mapping ϕ induces a CR equivalence between the natural universal coverings of M, M', which are the path spaces mod homotopic relation respectively over M, M' with the natural CR structure. This completes the proof. ☐

REFERENCES

[Bo] A. Boggess, *CR functions and the tangential Cauchy-Riemann complex*, CRC Press, Boca Raton, 1991

[CJ] S.-S. Chern and S. Ji, On the Riemann mapping theorem, *Ann. of Math.* **144**(1996) 421-439.

[Kr] S. G. Krantz, *Function theory of several complex variables*, Wadsworth, Inc., Belmont, 1992

[Pa1] W. K. Park, Normal forms of real hypersurfaces with nondegenerate Levi form, *http://xxx.lanl.gov/math.CV/9902034*

[Pa2] W. K. Park, Umbilic points and real hyperquadrics, *http://xxx.lanl.gov/math.CV/9902035*

[Pa3] W. K. Park, Analytic continuation of a biholomorphic mapping, *http://xxx.lanl.gov/math.CV/9902036*

[Pi] S. I. Pinchuk, On proper holomorphic mappings of strictly pseudoconvex domains, *Sib. Math. J.* **15**(1974) 644-649

[Ra] R. M. Range, *Holomorphic functions and integral representations in several complex variables*, Springer-Verlag, Berlin, 1986